

JAN 16 '53

# HYDROBIOLOGIA

ACTA HYDROBIOLOGICA, LIMNOLOGICA ET  
PROTISTOLOGICA

## EDITORES:

Gunnar Alm Drottningholm	H. d'Ancona Padova	Kaj Berg København
E. Fauré-Fremiet Paris	F. E. Fritsch London	H. Järnefelt Helsinki
P. van Oye Gent	K. Ström Oslo	W. R. Taylor Ann Arbor
N. Wibaut-Isebree Moens Amsterdam		





**HYDROBIOLOGIA** publishes original articles in the field of Hydrobiology, Limnology and Protistology. It will include investigations in the field of marine and freshwater Zoo- and Phytobiology, embracing also research on the Systematics and Taxonomy of the groups covered. Preliminary notices, polemics, and articles published elsewhere will not be accepted. The journal, however, contains reviews of recent books and papers.

Four numbers of the journal are published every year. Each number averages about 100 pages. Contributions must be clearly and concisely composed. They must be submitted in grammatically correct English, French, German, Italian or Spanish. Long historical introductions are not accepted. Protocols should be limited. Names of animals and plants must be given according to the laws of binominal nomenclature adopted at the recent International Congresses of Zoology and of Botany, including the author's name; it is desirable that the latter should be given in full. Measures and weights should be given in the decimal system. Every paper has to be accompanied by a short summary, and by a second one, written in an alternative language.

Manuscripts should be typewritten in double spacing on one side of the paper. The original should be sent. Original drawings should be submitted. Text figures will be reproduced by line engraving and hence should not include any shading, although figures which cannot be reproduced in this manner will be accepted if necessary. All drawings should be made on separate sheets of white paper, the reduction desired should be clearly indicated on the margin. The approximate position of text-figures should be indicated on the manuscript. A condensed title, should be cited as follows: in the text — AHLSTROM (1934); in the references — AHLSTROM, E. H., 1934. Rotatoria of Florida; *Trans. Amer. Micr. Soc.* 53: 252—266. In the case of a book in the text — HARVEY (1945); in the references — HARVEY, H. W.: Recent Advances in the Chemistry and Biology of Sea Water, Cambridge Univ. Pr., London 1945. Author's names are to be marked for printing in small capitals, latin names of animals and plants should be underlined to be printed in italics.

The various types of printing should be indicated by underlining the words in the following way:

- |           |   |
|-----------|---|
| =====     | CAPITALS, e.g. for headlines; preferably <i>not</i> in the text.  |
| ~~~~~     | or straight blue line: SMALL CAPITALS, e.g. <i>all</i> names of persons, both in the text and in the references.            |
| —————     | <b>heavy type</b> , e.g. for sub-titles; preferably <i>not</i> in the text.   |
| ~~~~~     | or straight red line: <i>italics</i> , e.g. <i>all</i> Latin names of plants and animals, except those in lists and tables. |
| - - - - - | spaced type.  |

Manuscripts may be sent to any member of the board of editors or directly to the *hon. secretary*, Prof. Dr. P. van Oye, 30, St. Lievenslaan, Ghent, Belgium, to whom proofs must be returned after being clearly corrected. Fifty free reprints of the paper with covers will be furnished by the publishers. Orders for additional copies should be noted on the form which is enclosed with the galleyproofs.

*Books and reprints are to be sent to the honorary secretary directly.*



# Cyclops Magnus from Amchitka, Alaska

GABRIEL W. COMITA

Dept. of Zoology <sup>1)</sup>

University of Washington

It is the intention of this paper to supply a small amount of data that may contribute to the clarification of the status of a cyclopoid copepod included in the viridis-vernalis group, *Cyclops magnus* Marsh 1920.

C. D. MARSH first established *C. magnus* from specimens found during the Canadian Arctic Expedition from areas in northern Canada. At a later date it was found at Churchill, Manitoba. It seems not to have been recorded from other nearby areas, perhaps due to a scarcity of collections from nearby regions and some probable mis-identifications by those few investigators who have collected in those areas.

The data presented here are taken from specimens collected on the island of Amchitka in the Aleutians. About two dozen specimens were taken from a small drainage pool being fed from tundra seepage water. The water appeared to have a considerable amount of vegetational leechings, being clear, but colored. The temperature of the water was approximately 45° F. and the water was constantly moving. Data taken from these specimens follow.

The measurements of length were (mm.):

	♀	♂
<i>nonovigerous</i>	<i>ovigerous</i>	
1.87	2.13	1.7
1.87	2.17	
1.87	2.4	
1.9	2.7 <sup>2)</sup>	
1.96		
2.10		
2.10		
2.10		
2.47		

<sup>1)</sup> The collection of specimens studied here was made possible through funds provided by the Office of Naval Research.

<sup>2)</sup> appeared abnormally extended.

These length measurements indicate an overall greater length than has its near relative, *C. viridis*, with which it can be easily confused.

The postero-lateral angles of the dorsa were without protrusions (see figure 3).

In every specimen examined the posterior border of the abdominal segments (figure 2) was coarsely serrated except the last (figure 4), which had its posterior margin armed with small spines.

The inner margins of the furcal rami were covered with hairs (figure 4). These fine hairs were rather difficult to discern because they seemed to be finer than usual and were not as easily recognizable as are those on the lateral margins of setae. These hairs were not regularly spaced as are the secondary hairs on setae but appeared to be outgrowths without arrangement.

The relative lengths of the terminal furcal setae (innermost to outermost) were:

a)	2.1	5.7	4.7	1.2
b)	1.9	5.0	4.9	1.0
c)	2.0	7.0	5.5	1.0

The ratios of length of furcal ramus to length of innermost terminal furcal seta expressed as  $\frac{\text{length of seta}}{\text{length of furca}}$  were:

1.05	1.16	1.13	1.21	1.09	1.05
1.05	1.025	1.05	1.11	1.20	1.17


This ratio takes on a new importance in the light of these measurements. The literature indicates that this is one of the few constant differences between this species and *C. viridis* as well as *C. gigas latipes* and *C. capillatus*.

The first antennae were of 17 segments, no exceptions were noted. The esthete on the distal end of segment 12 of the antenna extended beyond the middle of segment 14 but not past its end.

The inner terminal spine of endopod 3 of the fourth swimming foot was shorter in these specimens (figure 1), having the following ratio to the outer terminal spine of the same segment:

$$\frac{\text{outer terminal spine}}{\text{inner terminal spine}} = 1.16$$

The terminal segment of the endopod of the fourth swimming foot itself had the following width: length quotient:  $\frac{\text{width}}{\text{length}} = 0.437$ ,



#### EXPLANATION OF PLATE

Figure 1. Terminal segment of endopod of  $P_4$ .

Figure 2. Ventral view of abdomen.

Figure 3. First four segments of metasome.

Figure 4. Furcal rami, ventral view.

Figure 5. Fifth foot.

All drawings are from the female, *C. magnus*.



being somewhat more elongate than the endopods on the specimens collected in the Canadian Arctic Expedition.

The armature on the swimming feet was the same as that for other described specimens of *C. magnus*; the spine formula being 2, 3, 3, 3.

The fifth foot was of the form shown in figure 5. In every case examined the tiny inner spine was not jointed to the distal segment of the fifth foot.

The shape of the seminal receptacle seemed not to be variable, having the anterior margin concave (figure 2) in those specimens examined.

## SUMMARY

About two dozen specimens of *Cyclops magnus* were taken from a tundra seepage pool on the island of Amchitka, Alaska. Measurements from these specimens are presented. These measurements contribute further evidence of the reliability of certain spine and setal ratios as a means for differentiating some members of the viridis-vernalis group.

## RIASSUNTO

Circa due dozzine di esemplari di *Cyclops magnus* furono raccolte da uno stagno (tundra seepage pool) sulla isola di Amchitka, Alaska. Misure di questi esemplari sono presentati. Queste misure contribuiscono evidenza addizionale che certe proporzioni spinose e setoli sono degne di fiducia per differenziare qualche membro del gruppo viridis-vernalis.

## REFERENCES

- GURNEY, R. - British Freshwater Copepoda, 3: 29—284. Ray Society, London 120, 1933.
- MARSH, C. D. - Report of the Canadian Arctic Expedition 1913—18. Crustacea. (J) Freshwater Copepoda. Report Canad. Arct. Exp. 7 (J): 1—25, 1920.
- YEATMAN, H. C. - American Cyclopoid Copepods of the viridis-vernalis Group, (Including a Description of *Cyclops carolinianus* n. sp.) Amer. Midland Naturalist 32, No. 1, pp. 1—90, 1944.

Address of the author:  
UNIVERSITY OF WASHINGTON  
Department of Zoology  
Seattle, 5 Washington

# Etudes consacrées aux animaux dulcicoles et recherches d'Hydrobiologie animale en France de 1945 à 1950<sup>1)</sup>

## 1. Poissons et Batraciens

par

MAURICE BLANC

Notre collègue P. BOURRELLY, Assistant au Laboratoire de Cryptogamie du Muséum National d'Histoire Naturelle de Paris, ayant donné dans ce même périodique un article bibliographique très complet <sup>2)</sup> portant sur les travaux d'algologie et de limnologie botanique effectués récemment en France, nous avons cru faire oeuvre utile en publiant, d'accord avec lui, une mise en point analogue relative aux travaux français d'hydrobiologie animale <sup>3)</sup>.

### POISSONS <sup>4)</sup>

En ce qui concerne les faunes et les ouvrages d'ensemble, signalons le petit manuel de P. GOSSOT (1946) relatif aux Poissons des eaux douces de France et destiné surtout aux amateurs, ainsi que le fascicule IV du Petit Atlas des Poissons (édit. Boubée) dont la parution avait été retardée par la mort du regretté J. PELLEGRIN et dont la réalisation est l'oeuvre de F. ANGEL (1946).

Parmi les articles bibliographiques, nous avons une excellente mise au point de CH. DEVILLERS (1950) sur la constitution et l'évolution du crâne chez les Poissons, un article très documenté de J. DAGET (1949) sur le rôle de la ligne latérale chez les Poissons et un

<sup>1)</sup> En ce qui concerne l'année 1950, seuls les travaux publiés au début de l'année sont mentionnés dans cet article qui a été rédigé en août 1950.

<sup>2)</sup> BOURRELLY P. Activités algologiques et limnologiques en France pendant les années 1947-1948. *Hydrobiologia*, 1949, II (no 1), pp. 72-83.

<sup>3)</sup> Nous ne citons ici que les travaux sur les Poissons d'eau douce et les Poissons amphibiotiques.



travail de CH. THIBAUT et O. THIBAUT sur l'adaptation chromatique chez les Poissons (1947).

En systématique, A. BARETS utilise les caractères biométriques pour distinguer plus sûrement certains Cyprinidés très proches l'un de l'autre, tels que *Cyprinus* et *Carassius* (1947) ou *Gardonus* et *Scardinius* (1948).

Au point de vue histologique et histochimique, les travaux sont assez nombreux pendant cette période. R. ARGAUD et J. K. GAN (1945) découvrent l'existence d'ilôts pancréatiques dans la paroi même de la vésicule biliaire de l'Anguille; ces pancréas minuscules possèdent à la fois les fonctions endocrine et exocrine. L. BERNER étudie la formation et la disposition des fibres collagènes qui relient la dent à son socle chez les Téléostéens (1945), puis il décrit la structure et le développement des dents chez *Charax puntazzo* Risso (1945) et chez *Gambusia affinis* Bd. Gd. (1947). C. BIMES décrit l'hépatopancréas des Poissons (1947). G. BOBIN (1949) précise la structure cytologique des corpuscules de Stannius (système interrénal postérieur) de l'Anguille et décrit l'évolution des lobules de la Civelle à l'adulte; il s'agit bien là d'une glande à sécrétion interne (système cortico-surrénal). G. KOBIS (1949) se livre à une étude histochimique du syncytium vitellin de l'alevin de *Salmo irideus* Gibb.; cette étude porte principalement sur le fer et sur les porphyrines ainsi que sur le stade d'apparition de l'hémoglobine. G. PALLOT, G. et J. TOURNEUX (1946) étudient les variations sexuelles de l'ilôt pancréatique des Téléostéens. Le Dr. L. THOMAS (1948) présente une thèse volumineuse et richement illustrée sur l'histopathologie des Poissons. Enfin, M. BLANC consacre une série de notes à l'étude de l'ostéogénèse chez les Téléostéens. Après avoir étudié la structure et le développement des arcs branchiaux, M. BLANC étudie en détail la structure des rayons des nageoires (1947) et interprète le tissu ostéoïde des Poissons comme un tissu osseux jeune susceptible de se transformer ensuite en tissu osseux véritable. Puis l'auteur se livre à des expériences de régénération osseuse sur des espèces variées; ces expériences de régénération donnent lieu à des observations morphologiques (1948, 1950), histologiques (1949) ou même physiologiques (1948, 1949), ces dernières étant généralement faites en collaboration avec Mme J. BUSER (1949).

En hématologie, L. ARVY et M. GABE (1948) étudient le vacuome des érythrocytes chez plusieurs Téléostéens d'eau douce. M. GABE et C. VEIL (1948) décrivent les basophiles tissulaires de l'écaille de la Carpe. Enfin, M. LAUR (1947, 1949) s'occupe particulièrement des leucocytes des Poissons.

L'embryologie des Poissons, qu'elle soit normale ou expérimen-



ale, est surtout représentée pendant cette période par CH. DEVILLERS et par TH. THOMOPOULOS. CH. DEVILLERS publie d'abord une série de notes sur la genèse de certains os du crâne chez les Salmonidés (1946, 1947 et 1948). Puis il présente une thèse très remarquée (1947) sur le rôle des organes sensoriels dans la formation du crâne dermique des Téléostéens; à l'aide d'expériences d'excisions et de greffes, l'auteur met en évidence le rôle morphogénétique des neuromastes. Après sa thèse, CH. DEVILLERS publie une autre série de notes relatives cette fois à la segmentation de l'oeuf de Truite (1948, 1950). TH. THOMOPOULOS se consacre à l'embryologie topographique de la thyroïde de *Salmo fario* L. et de *Salmo irideus* Gibb. (1948, 1950) et suggère des vues tout-à-fait nouvelles sur le développement de cette glande qui, chez les Poissons, est fort aberrante.

Au point de vue de la reproduction des Poissons, citons un article de L. BERTIN (1947) sur les nids des Poissons, un article de Y. FRANÇOIS (1946) sur les principaux aspects de la reproduction chez les Poissons et deux notes de G. BRESSE (1946) sur la reproduction chez les Bouvières. L. GALLIEN publie plusieurs notes (1945, 1946 et 1947) sur la reproduction et la génétique de *Lebistes reticulatus* Peters.

En physiologie, les travaux sont nombreux. D. ABBE—FESSARD, C. CHAGAS, A. COUCEIRO et A. FESSARD (1948) étudient la production d'électricité chez les Gymnotes. Mme J. BUSER, soit seule (1948, 1950), soit en collaboration avec M. BLANC (1949), étudie la biochimie et la physiologie de l'ostéogénèse chez les Téléostéens. P. BUSER (1949) publie une note de physiologie nerveuse sur l'excitation du nerf optique chez le Poisson-Chat et compare cette expérience avec une expérience analogue faite chez la Grenouille, R. G. BUSNEL, A. DRILHON et A. RAFFY étudient la physiologie générale des Poissons de la famille des Salmonidés (1946). M. LAFFON (1947) recherche la façon dont les réserves vitellines sont utilisées au cours du développement chez les alevins de Truite. R. LHOTTE (1945) publie une note relative à l'influence de la salinité sur le fonctionnement du coeur chez la Civelles. CH. THIBAUT étudie de façon très approfondie le comportement des Poissons, et plus particulièrement de la Carpe, vis à vis de la lumière (1946, 1947, et 1949) et leurs réactions aux variations d'éclairement. V. VILTER publie une série de notes sur divers aspects de la métamorphose chez l'Anguille; il montre notamment que l'hypophyse commande les modifications pigmentaires accompagnant la métamorphose (1945), que la brusque augmentation de la croissance stomacale des Civelles est due au passage dans l'eau douce (1945) et que la thyroxine ne semble pas intervenir sur le déclenchement de la métamorphose de l'Anguille contrairement à ce qui se passe chez les Batraciens (1946). Pour terminer ce

paragraphe sur la physiologie ichthyologique, il faut faire une mention spéciale des travaux de M. FONTAINE qui, avec l'aide de ses élèves, notamment O. CALLAMAND, J. LELOUP et M. OLIVEREAU, consacre une grande partie de son activité depuis plusieurs années à l'étude du problème des migrations chez les Poissons et notamment à l'étude du rôle des différents facteurs, externes et internes, dans ces migrations compliquées (voir nombreuses références de 1945 à 1950).

Mais les Poissons des eaux douces métropolitaines ne sont pas les seuls à être étudiés par les Français, car depuis quelque temps les espèces exotiques d'eau douce sont également à l'honneur. C'est ainsi que L. BERTIN (1948) consacre un article important à la biogéographie des Poissons d'eau douce de Madagascar, et que son assistante, R. ESTEVE, publie une note sur les barbeaux marocains (1947) et une autre note sur les Poissons du Sahara Central (1949). En Afrique Occidentale, l'étude des poissons d'eau douce est principalement entreprise par l'Institut Français d'Afrique Noire (directeur TH. MONOD) qui possède, d'une part un laboratoire lagunaire à Abidjan, Côte d'Ivoire (directeur J. L. TOURNIER), avec un laboratoire flottant automoteur, le Baléo II, et d'autre part une Station hydrobiologique en construction, à Diafarabé sur le Niger, Soudan (directeur: J. DAGET), avec des bassins d'élevage et un chaland-laboratoire à moteur: le Tilapia. Alors qu'Abidjan a consacré jusqu'ici son activité principale aux problèmes hydrologiques, Diafarabé a pour objectif essentiel l'étude des poissons (faunistique, biologie, développement, biométrie, scalimétrie, ichthyologie appliquée). TH. MONOD consacre de plus un numéro entier de *Cybiu*m (no 4, 1949) aux travaux sur les eaux douces africaines et publie (1945, 1947, 1949 et 1950), ainsi que J. DAGET (1948, 1949 et 1950), plusieurs notes sur les Poissons de ces régions. J. DAGET publie d'autre part une thèse importante sur les Polyptéridés (1949). Rappelons également que P. BUDKER, dans son livre sur „la vie des requins” (1947) consacre un important chapitre aux requins d'eau douce, et que J. CHAUX se spécialise dans l'étude des représentants de la famille des Siluridés (1948, 1949). Enfin, pour finir, signalons la belle remise en marche, grâce aux efforts de G. GOUSSEF, de l'Aquarium du Musée des Colonies dont l'activité avait été presque entièrement stoppée par suite de la guerre.

Avant d'en terminer avec les études sur les Poissons d'eau douce, il nous reste à évoquer les travaux proprement piscicoles. Ces travaux, qui sont poursuivis dans plusieurs centres, ont été, pendant la période considérée, l'objet d'encouragements particulièrement attentifs de la part de l'Administration des Eaux et Forêts et des services qui en dépendent ou collaborent avec elle.



L'élevage des Salmonidés constitue une des principales branches de la pisciculture en France. R. LIGNIERES et H. HOESTLANDT (1949, 1950) examinent le problème délicat de l'alimentation artificielle des jeunes alevins de *Salmo irideus* Gibb. R. TAGAND, H. HOESTLANDT et L. EYRAUD (1947) étudient les principaux facteurs de croissance de cette même espèce. Enfin, R. VIBERT s'occupe particulièrement de la répartition des Truites et des Saumons dans les conditions naturelles (1945, 1948, 1949 et 1950).

L'élevage des Cyprinidés, et principalement des Carpes, donne lieu également à des recherches importantes. R. CHIMITS et S. KRECZMER (1946, 1947) s'occupent de l'élevage des alevins de Carpe dans différentes conditions et M. LEFEVRE (1948) étudie la nutrition de la Carpe en étang.

L'élevage des alevins de Brochet ainsi que le repeuplement des eaux douces en Brochets commencent à faire l'objet de recherches assez approfondies de la part de quelques auteurs, principalement de P. CHIMITS (1947) et de J. PORTAL (1947).

L'Alose du Rhône constitue un matériel précieux abondamment utilisé par C. GALLOIS (1946, 1947) qui réussit même à pratiquer la fécondation chez cette espèce, de même que l'abbé H. HOESTLANDT (1947, 1948). Le développement de la Perche est étudié par J. ARLET (1945). Enfin P. CHIMITS (1947), déjà cité, s'occupe également de l'acclimatation de *Gambusia holbrooki* dans la région des Landes.

Pour terminer ce chapitre, signalons le petit livre de M. BOUCHER (1948) intitulé "l'aquarium équilibré", dans lequel les amateurs aquariophiles trouveront tous les renseignements techniques nécessaires sur l'élevage en appartement des Poissons d'eau froide et d'eau chaude.

## BATRACIENS

En ce qui concerne la systématique des Batraciens, signalons principalement les travaux de F. ANGEL et ceux de J. GUIBÉ. F. ANGEL publie le "Petit Atlas des Amphibiens et des Reptiles", en deux fascicules (1946), puis la Faune de France des Reptiles et des Batraciens (1946) et enfin un traité plus général intitulé "Vie et Mœurs des Amphibiens" (1947). D'autre part, il publie en collaboration avec M. LAMOTTE une note sur la biologie d'un Crapaud vivipare du genre *Nectophrynoïdes*. Avec J. GUIBÉ il publie quelques notes de pure systématique (1945). J. GUIBÉ publie en outre seul plusieurs notes de systématique (1945, 1947), ainsi qu'un court article sur la biogéographie des Batraciens de Madagascar (1948). Dans le même ordre d'idées, P. REY démontre l'existence de races

géographiques chez *Bufo vulgaris* Laur. (1946) et L. VALLEE décrit les races sexuelles françaises de *Rana temporaria* L. (1947).

Du point de vue anatomique et histologique, signalons le travail de R. COUTEAUX (1945) sur les rapports entre nerfs et muscles chez la Grenouille, et celui de F. MARCEAU (1946) sur la terminaison du pneumogastrique dans le muscle cardiaque de la Grenouille et du Crapaud. M. LAUR (1948) présente une note sur l'hématologie du Triton.

L'embryologie des Batraciens donne lieu à des travaux importants. Ce sont en effet ceux des Vertébrés dont les oeufs et les embryons se prêtent le mieux à l'expérimentation. P. ANCEL et P. VINTEMBERGER, dans une série de notes (1947, 1948 et 1949), étudient la segmentation de l'oeuf. D. BOGORAZE (1946) se livre à des recherches sur le système nerveux des têtards d'Anoures. R. CAMBAR publie une douzaine de notes (1947, 1948 et 1949) sur le système rénal; il étudie tour à tour la formation du pronéphros, puis du mésonéphros chez les Anoures et recherche les éléments inducteurs du système rénal. P. HATT (1945) étudie la blastotomie chez le Triton. R. LALLIER (1949) étudie l'action de différentes substances sur la gastrulation. J. ROSTAND, après avoir examiné en collaboration avec H. MUGARD (1946) l'action du gaz carbonique sur les oeufs de Batraciens, cherche à perturber le processus normal du développement de l'oeuf (1946, 1947, 1948 et 1949). P. SENTIN (1948, 1949) s'occupe de la mitose et de l'action des substances antimitotiques sur l'oeuf de Batraciens. R. STOLL (1946) décrit les premiers stades de la segmentation de l'oeuf chez *Bufo vulgaris* Laur. Enfin, J. A. THOMAS, J. ROSTAND et J. GREGOIRE (1946) mettent en évidence l'action inhibitrice de la ribonucléase sur la segmentation des oeufs de Grenouille.

En Physiologie, les travaux sont nombreux également, la Grenouille étant, comme chacun sait, un parfait animal de laboratoire. L. ARVY, J. A. BOIFFARD et M. GABE (1947) étudient l'action des rayons X sur l'hypophyse de *Rana temporaria* L., puis les modifications hématologiques provoquées sur les sujets ainsi irradiés. L. ARVY décrit un dimorphisme sexuel sanguin chez *Rana temporaria* L. et chez *Bufo vulgaris* Laur. (1947), puis étudie l'action de la glande thyroïde sur l'hématopoïèse chez la Grenouille (1949, 1950). C. CHAMPY et M. DEMAY recherchent le rôle du système nerveux dans le développement des callosités (1949). Les mêmes auteurs décrivent les dispositifs d'adaptation au saut chez les Grenouilles (1949), J. CREZE (1949) présente deux petites notes sur le déterminisme de la ponte chez *Xenopus laevis*. M. DELSOL (1950) cherche à évaluer l'activité de la thyroïde chez les Têtards de Discoglosses à l'aide du thiouracile. M. GABE (1946) se livre à des expériences d'autogreffe hépatique sur la grenouille d'hiver. L. GALLIEN étu-



die la callosité de *Xenopus laevis* (1947), puis l'action de la prégnénolone comme antithyroïdien chez le têtard de Discoglosse (1949) et l'action de l'éthinyl-testostérone sur le développement de *Rana temporaria* L. (1949). C. GAUTHIER dans une courte note (1946) étudie la naissance du réflexe amoteur chez le Crapaud. J. JOLLY étudie la disposition et la formation du système lymphatique des Batraciens (1945, 1946). Ce même auteur étudie également quelques phénomènes de régénération (1946, 1949) chez ces animaux. M. LECAMP étudie les phénomènes de régénération chez les Batraciens, notamment l'action de diverses substances sur la régénération, ainsi que la culture des tissus de régénération in vitro (1947, 1948, 1949). J. LHOSTE et P. ROTH (1946) étudient le développement des oeufs de Grenouille. D. PICARD et H. TUCHMANN—DUPLESSIS (1946) étudient le rôle du système réticulo-endothélial chez le Triton alpestre. A. RAFFY donne une revue intéressante et documentée sur la physiologie des animaux amphibies; des exemples pris chez les Batraciens y sont souvent cités. P. ROTH publie une thèse (1945) et de nombreuses notes (1946, 1947, 1948 et 1950) sur l'action de diverses substances hormonales et notamment de la thyroxine sur la métamorphose des Batraciens. A. SERFATY (1946) donne une courte note sur la résistance des Grenouilles à l'asphyxie. F. STUTINSKY (1945, 1946 et 1947) étudie de façon suivie la mélanocinèse des Batraciens et l'action du lobe intermédiaire de l'hypophyse sur celle-ci. J. A. THOMAS et I. BORDERIOUX (1947, 1948) réalisent de belles expériences de survie avec des fragments d'organes de Triton conservés dans un milieu spécialement préparé. H. TUCHMANN—DUPLESSIS travaille sur les glandes endocrines (thyroïde, hypophyse et gonades) du Triton, étudie leurs variations saisonnières, leurs interactions et leur influence sur la régénération (1945, 1946, 1948 et 1949). Enfin, J. TUSQUES étudie le rôle de la thyroxine sur le développement des têtards (1949).

Pour terminer ce chapitre, il nous reste à citer quelques publications à tendance écologique, telles que celle de E. BRUMPT (1945) à propos de l'utilisation de *Xenopus laevis* dans la lutte contre les larves de Moustiques, celle de Y. MORON (1947) sur l'écologie, la biologie et l'anatomie de *Xenopus tropicalis* rapporté de Haute-Guinée par M. LAMOTTE, et celle de R. PAULIAN et A. VILARDEBO (1946) sur l'alimentation des Batraciens en Basse Côte d'Ivoire.

(Laboratoire des Pêches Coloniales du Museum National d'Histoire Naturelle de Paris et Laboratoire d'Hydrobiologie du Centre National de la Recherche Scientifique à Gif-sur-Yvette - Seine et Oise).

## ADDENDUM

*Travaux parus en 1950.*

### POISSONS<sup>1)</sup>

En systématique, L. BERTIN publie un mémoire sur la curieuse collection de Poissons en herbier du naturaliste MICHEL ADANSON et sur la façon dont celui-ci concevait la classification ichthyologique.

En ce qui concerne les travaux d'ordre physiologique, nous pouvons signaler plusieurs notes ou articles de M. FONTAINE et de ses collaborateurs sur le Saumon et les principales variations biochimiques observées chez cet animal aux différentes périodes de sa vie. J. BARON publie plusieurs notes sur l'étude de l'équilibre chez les Poissons et plus particulièrement sur le rôle des muscles moteurs oculaires dans l'équilibration. L'abbé H. HOESTLANDT effectue, en collaboration avec H. VINCENT, des essais relatifs à l'action de l'acide 2—4 dichlorophénoxyacétique sur les Poissons et les Batraciens. A. JULLIEN, P. LAURENT et J. RIPPLINGER (1950) effectuent des dosages d'acétylcholine chez plusieurs Poissons d'eau douce. Ils réalisent chez les Mollusques et les Poissons la préparation sur le vivant de la branche cardiaque du pneumogastrique et pratiquent des excitations tout en contrôlant le rythme du coeur par des enregistrements graphiques.

En histologie, TH. THOMOPOULOS étudie la thyroïde de *Gymnarchus niloticus* Cuv. et fait des comparaisons avec les Sélaciens et les Salmonidés.

Au point de vue de la reproduction, G. BRESSE donne deux nouvelles notes sur la curieuse reproduction des Bouvières et le rôle des Moules d'eau douce au cours de celle-ci.

Les espèces exotiques donnent lieu également à des travaux intéressants. TH. MONOD publie de nouvelles observations sur les Poissons d'eau douce de l'A.O.F., tandis que J. DAGET poursuit ses travaux à la Station Hydrobiologique de Diafarabé. D'autre part, J. DURAND présente un important mémoire sur l'hématologie des Poissons d'Indochine. Enfin, J. PUYO fait paraître une faune générale des Poissons de la Guyane française, dont il rassemblait les éléments depuis de nombreuses années.

Pour terminer, signalons qu'au point de vue proprement piscicole, R. VIBERT donne une nouvelle étude du Saumon de l'Adour, tandis que J. WURTZ-ARLET présente quelques observations sur la Vandoise.

<sup>1)</sup> Nous envisageons uniquement les Poissons d'eau douce ou amphibiotiques.



## BATRACIENS

En systématique, L. BERTIN assure la publication à titre posthume de deux notes de F. ANGEL, décédé dans l'année; l'une de ces notes concerne les Batraciens de Madagascar et l'autre ceux des Monts Nimba (Guinée). D'autre part, J. GUIBE étudie lui aussi les Batraciens de Madagascar, soit en collaboration avec J. MILLOT, soit seul.

En ce qui concerne les travaux de physiologie, L. GALLIEN, dans un important mémoire, étudie la différenciation du sexe chez le Discoglosse et la Grenouille rousse, et P. ROTH continue la série de ses publications relatives à l'action des hormones sur le développement et les métamorphoses de l'Axolotl. Rappelons que H. HOESTLANDT et H. VINCENT essayent l'action de l'acide 2—4 dichlorophénoxyacétique sur les Batraciens Anoures en même temps que sur les Poissons, ainsi que nous l'avons signalé plus haut. Enfin H. TUCHMANN—DUPLESSIS étudie l'action inhibitrice des sels de béryllium sur les phénomènes de régénération chez différentes espèces de Tritons.

Au point de vue cytologique, L. GALLIEN et H. MUGARD font une démonstration technique d'imprégnation argentique et de coloration au Feulgen de l'extrémité caudale des larves de Pleurodèles hétéroplôides.

Enfin, en neurologie, M. JACQUOT publie deux notes sur le développement cérébral des Batraciens Urodèles.

(Laboratoire des Pêches Coloniales du Muséum National d'Histoire Naturelle de Paris et Laboratoire d'Hydrobiologie du C.N.R.S. à Gif-sur-Yvette - Seine et Oise).

## BIBLIOGRAPHIE

### POISSONS

- ABBE-FESSARD, D., CHAGAS, C., COUCERIO, A. et FESSARD, A. - Le problème de la propagation de l'onde de décharge dans l'organe électrique du Gymnote (*Electrophorus electricus*). *C. R. Acad. Sci. Paris*, 1948, 226, pp. 1309—1310.
- Hypothèse sur le mécanisme de la propagation de la décharge chez le Gymnote (*Electrophorus electricus*). *C. R. Acad. Sci. Paris*, 1948, 226, pp. 1474—1476.
- ANGEL, F. - Petit Atlas des Poissons, fascicule 4: Poissons des eaux douces; espèces exotiques et d'ornement. N. Boubée, Paris, 1946.
- ARGAUD, R. et GAN, J. K. - Sur l'inclusion de minuscules pancréas accessoires dans l'épaisseur même de la paroi de la vésicule biliaire chez *Anguilla vulgaris* Cuv. *Bull. Mus. Nat. Hist. Nat. Paris*, 1945, 17, pp. 466—470.
- ARLET, J. - Note sur la fécondation artificielle et le développement des alevins de Perche. *Bull. Fr. Piscicult.*, 1945, 138, pp. 32—37.

- ARVY, L. et GABE, M. - Les corps de Golgi des érythrocytes chez quelques Téléostéens d'eau douce. *C. R. Soc. Biol.*, 1948, 142, pp. 278—279.
- BARETS, A. - Essai d'analyse biométrique des genres *Cyprinus* et *Carassius*. *Bull. Mus. Nat. Hist. Nat.* 1947, 19, pp. 70—73.
- Sur un indice numérique utile à la détermination de *Gardonus rutilus* (L.) et *Scardinius erythrophthalmus* (L.). *Bull. Mus. Nat. Hist. Nat. Paris*, 1948, 20, pp. 72—74.
- BARON, J. - Contribution à l'étude de l'équilibre chez les Poissons. *Bull. Soc. Zool. Fr.*, 1950, 75, pp. 247—252.
- Musculature extrinsèque et équilibre des Poissons. *C. R. Acad. Sci. Paris*, 1950, 230, p. 2231.
- Relations entre les muscles moteurs oculaires, les nageoires et l'équilibre des Poissons. *C. R. Acad. Sci. Paris*, 1950, 231, pp. 1087—1088.
- BERNER, L. - Le ligament dentaire des Téléostéens. *Bull. Soc. Zool. Fr.*, 1945, 70, pp. 26—27.
- L'organe prédentaire de *Charax puntazzo* Risso. *Bull. Soc. Zool. Fr.*, 1945, 70, pp. 156—158.
- Le développement dentaire chez *Gambusia affinis* (Baird et Girard). *Bull. Soc. Zool. Fr.*, 1947, 72, pp. 22—30.
- BERTIN, L. - Les nids des Poissons. *Feuille Nat.*, 1947, 2, p. 22.
- Considérations biogéographiques sur les Poissons d'eau douce de Madagascar. *Mém. Inst. Sci. Madagascar*, 1948, série A, 1, pp. 169—179.
- Les Poissons en herbier et le système ichthyologique de MICHEL ADANSON. *Mém. Mus. Nat. Hist. Nat.*, 1950, A, 1, 45 p.
- BIMES, C. - L'hépatopancréas des Poissons. *Arch. Anat. Hist. Embryol.*, 1947, 30, pp. 53—67.
- BLANC, M. - Etude comparative de l'ossification des arcs branchiaux chez les Poissons Téléostéens. Diplôme d'Etudes Supérieures, Fac. Sci. Paris, no d'ordre 902, 20 Novembre 1944, 20 p. dactylographiées, 11 fig.
- L'ossification des arcs branchiaux chez les Poissons Téléostéens. *Bull. Soc. Zool. Fr.*, 1944, 69, pp. 226—230, 1 fig.
- Structure histologique des rayons de nageoires chez les Poissons Téléostéens. *Bull. Soc. Zool. Fr.*, 1947, 72, pp. 17—22.
- Sur la réparation des nageoires de Poissons Téléostéens après amputation d'un volet médian. *Arch. Zool. Exp. Gén.*, 1948, 85, N. et Rev. no 4, pp. 184—188.
- Sur la destinée des fragments de rayons osseux isolés de leur base dans les nageoires de Poissons Téléostéens. *C. R. Acad. Sci. Paris*, 1948, 226, pp. 1466—1468.
- Etude histologique de la régénération des nageoires chez quelques Poissons Téléostéens. *Arch. Anat. Microsc. Morph. Exp.*, 1949, 38, pp. 52—54, 4 fig.
- et BUSER, J. - Action de la température sur l'ostéogénèse réparatrice chez les Poissons Téléostéens. *Bull. Soc. Zool. Fr.*, 1949, 74, pp. 167—170.
- BLANC, M. - Convergence morphologique des épines des nageoires de Labridés et de Poissons-Chats. *Feuille Nat.*, 1950, 5, pp. 11—12, 2 fig.
- BOBIN, G. - Images histo-cytologiques des corpuscules de Stannius de l'Anguille Européenne. *Arch. Zool. Exp. Gén.*, 1949, 86, pp. 1—7.
- BOUCHER, M. - L'aquarium équilibré. Paris (2ème édition), 1948, 74 p.



- BRESSE, G. - La reproduction des Bouvières. *Bull. Soc. Zool. Fr.*, 1946, 71, pp. 115—117.
- La reproduction des Bouvières. *Bull. Fr. Piscicult.*, 1946, 141, pp. 177—187.
- G. - Rôle des Moules d'eau douce dans la reproduction des Bouvières. *Bull. Fr. Piscicult.*, 1950, 159, pp. 47—52.
- Rôle des Moules d'eau douce dans la reproduction des Bouvières. *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, pp. 333—335.
- BUDKER, P. - La vie des requins. Gallimard, Paris, 1947, 277 p. (voir le chapitre VI: Requins d'eau douce, pp. 155—169).
- BUSER, J. - Modifications chimiques locales au cours de la régénération osseuse des nageoires de Poisson-Chat. *C. R. Soc. Biol.*, 1948, 142, pp. 799—800.
- et BLANC, M. - Action de la lumière sur l'ostéogénèse réparatrice chez le Poisson-Chat. *Bull. Soc. Zool. Fr.*, 1949, 74, pp. 170—172.
- et ——— Etude des variations du degré d'ossification des différentes parties de la nageoire pectorale du Poisson-Chat. *Arch. Zool. Exp. Gén.*, 1949, 86, N. et Rev. no 1, pp. 8—12, 2 fig.
- BUSER, J. - La prégnéninone utilisée comme facteur antithyroïdien et son rôle inhibiteur sur la régénération chez *Lebistes reticulatus* et *Ameiurus nebulosus*. *Bull. Soc. Zool. Fr.*, 1950, 75, pp. 75—80, 3 fig.
- BUSER, P. - Analyse de la réponse mésencéphalique à la stimulation du nerf optique chez le Poisson-Chat. *C. R. Soc. Biol.*, 1949, 143, pp. 817—819.
- BUSNEL, R. G., DRILHON, A., RAFFY, A. - Recherches sur la physiologie des Salmonidés. *Bull. Inst. Océanogr. Monaco*, 1946, 893, pp. 1—23.
- CALLAMAND, O. - Observations sur l'activité hématopoïétique des foies de quelques Poissons. *Bull. Soc. Centr. Aqu.*, 1947, 54, pp. 13—17.
- Aptitude de l'Anguille à former des antitoxines. Rapports possibles avec les mystères de sa reproduction. *Bull. Mus. Nat. Hist. Nat. Paris*, 1949, 21, pp. 512—515.
- CHAUX, J. - Note sur *Cryptopterus bicirrhis* C. V. *Bull. Mus. Nat. Hist. Nat. Paris*, 1948, 20, pp. 502—507.
- et †FANG, P. W. - Catalogue des Siluroïdes d'Indochine de la collection du Laboratoire des Pêches Coloniales au Muséum, avec la description de six nouvelles espèces. *Bull. Mus. Nat. Hist. Nat. Paris*, 1949, 21, pp. 194—201 et pp. 342—346.
- CHIMITZ, P. et KRECZMER, S. - Note sur la méthode d'élevage de l'alevin de Carpe en frayères Dubisch et bassin de grossissement. *Bull. Fr. Piscicult.*, 1946, 142, pp. 30—39.
- Note sur la méthode d'élevage de l'alevin de Carpe en frayères Dubisch et bassin de grossissement (suite et fin). *Bull. Fr. Piscicult.*, 1947, 144, pp. 137—143.
- CHIMITZ, P. - Note sur le repeuplement artificiel du Brochet. *Bull. Fr. Piscicult.*, 1947, 146, pp. 16—24.
- Note sur l'acclimatation du *Gambusia holbrooki* dans les étangs des Landes. *Bull. Fr. Piscicult.*, 1947, 147, pp. 79—82.
- DAGET, J. - La collection de poissons d'eau douce de l'I. F. A. N. *Catalogue III de l'Institut Français d'Afrique Noire*, Dakar, 1948, 59 p., 14 fig.
- Note sur les *Polypterus* du Niger moyen, notamment ceux de l'espèce *senegalus*. *Bull. Mus. Nat. Hist. Nat. Paris*, 1948, 20, pp. 82—86.
- Les *Synodontis* (Siluridés) à polarité pigmentaire inversée. *Bull. Mus. Nat. Hist. Nat. Paris*, 1948, 20, pp. 239—423.

- La ligne latérale: ses fonctions. *L'Année Biologique*, 1949, 25, pp. 305—322.
- Note préliminaire sur les affinités phylogénétiques des Polyptéridés. *Bull. Mus. Nat. Hist. Nat. Paris*, 1949, 21, pp. 532—536.
- Révision des affinités phylogénétiques des Polyptéridés. Thèse, Paris, 1949. (*Mémoires de l'I. F. A. N.*, Dakar, 1950, 11, 178 p.).
- Le Tinéni: poisson migrateur des eaux douces africaines. *Cybium*, 1949, 4, pp. 62—67.
- Poissons d'eau douce de Sierra Leone. *Notes Africaines (Bull. Inform. Corr. I. F. A. N.)*, 1950, 46, pp. 55—57.
- Poissons d'eau douce de la région côtière du Togo et du Dahomey. *Notes Africaines (Bull. Inform. Corr. I. F. A. N.)*, 1950, 46, pp. 57—59.
- La passe à poissons de Markala. *Bull. I. F. A. N.*, 1950, 12 (sous presse)
- Poissons. in: Contribution à l'étude de la Réserve Naturelle Intégrale des Monts Nimba. (sous presse)
- DEVILLERS, CH. - Etude expérimentale du rôle morphogénétique des fossettes sensorielles (pit-organs) de *Salmo fario*. *C. R. Acad. Sci. Paris*, 1946, 223, pp. 1180—1181.
- Explantations in vitro de blastodermes de Poissons (*Salmo*, *Esox*). *Experientia*, 1947, 3, p. 71.
- Recherches sur le crâne dermique des Téléostéens. *Ann. Paléontol.*, 1947, 33, pp. 1—94, 6 pl.
- La genèse des organes sensoriels latéraux de la Truite (*Salmo fario* et *Salmo irideus*). *C. R. Acad. Sci. Paris*, 1948, 226, pp. 334—356.
- Mécanisme gastruléenne de l'oeuf de Truite. *C. R. Acad. Sci. Paris*, 1948, 226, pp. 1310—1312.
- Suppression du matériel chordal dans la gastrula de l'oeuf. *C. R. Acad. Sci. Paris*, 1948, 227, pp. 1411—1413.
- Le cortex de l'oeuf de Truite. *Ann. Stat. Centr. Hydrob. Appl.*, 1948, 2, pp. 229—249.
- Action des solutions hypertoniques sur le développement de l'oeuf de Truite. *C. R. Acad. Sci. Paris*, 1950, 230, pp. 471—472.
- Quelques aspects de l'évolution du crâne chez les Poissons. *Année Biologique*, 1950, 26, 5, pp. 145—180, 17 fig.
- DURAND, J. - Etude morphologique du sang, de l'immunité naturelle et acquise, chez quelques Poissons Indochinois (marins et d'eau douce). *Ann. Inst. Ocean. Monaco*, 1950, 25, pp. 109—206.
- ESTÈVE, R. - Etude biométrique des Barbeaux marocains. *Bull. Mus. Nat. Hist. Nat. Paris*, 1947, 19, pp. 265—270.
- Poissons du Sahara Central. *Bull. Soc. Zool. Fr.*, 1949, 74, pp. 19—20.
- FONTAINE, M., DELATTRE, S. et CALLAMAND, O. - Influence des variations de salinité sur la teneur en hématies de deux Téléostéens (*Anguilla anguilla* L. et *Cyprinus carpio* L.). *Bull. Inst. Océan. Monaco*, 1945, 886, pp. 1—11.
- FONTAINE, M. - Vues actuelles sur les migrations des Poissons. *Experientia*, 1946, pp. 233—237.
- et CALLAMAND, O. - Sur les modifications du milieu intérieur des Poissons au cours des changements de salinité et leur interprétation. *C. R. Acad. Sci. Paris*, 1946, 222, pp. 198—200.
- et LELOUP, J. - Action d'antithyroïdiens (aminothiazol et phénylthiourée) sur la cuprémie des Vertébrés poecilothermes. *C. R. Soc. Biol.*, 1947, 141, pp. 148—149.
- et OLIVIEREAU, M. - La glande thyroïde du Saumon Atlantique (*Salmo*



- salar L.) ♀ au cours de sa vie en eau douce. *C. R. Acad. Sci. Paris*, 1947, 224, pp. 1660—1662.
- et CALLAMAND, O. - Sur l'activité antigonadotrope du sérum d'Anguille. *C. R. Acad. Sci. Paris*, 1947, 225, pp. 143—145.
- FONTAINE, M. - Physiologie du Saumon. *Ann. Stat. Centr. Hydrob. Appl.*, 1948, 2, pp. 153—183.
- , LACHIER, F., LELOUP, J. et OLIVEREAU, M. - La fonction thyroïdienne du Saumon (*Salmo salar* L.) au cours de sa migration reproductrice. *J. Physiol.*, 1948, 40, p. 182.
- et CALLAMAND, O. - Nouvelles recherches sur le déterminisme physiologique de l'avalaison des Poissons migrateurs amphibiotes. *Bull. Mus. Nat. Hist. Nat. Paris*, 1948, 20, pp. 317—230.
- FONTAINE, M. - Les migrations des Vertébrés envisagées au point de vue de la pathologie comparée. *Rev. Pathol. Comp.*, 1949, 49, 603, pp. 10—15.
- et CALLAMAND, O. - La lipase sérique chez un Cyclostome (*Petromyzon marinus* L.) et divers Poissons Téléostéens. *Bull. Inst. Océan. Monaco*, 1949, 943, pp. 1—5.
- La lipase sérique du Saumon (*Salmo salar* L.) à diverses étapes de son développement et de ses migrations. *Bull. Inst. Océan. Monaco*, 1949, 944, pp. 1—2.
- et OLIVEREAU, M. - Hypophyse et euryhalinité chez l'Anguille. *C. R. Acad. Sci. Paris*, 1949, 228, p. 513.
- FONTAINE, M. et OLIVEREAU, M. - L'hypophyse du Saumon (*Salmo salar* L.) à diverses étapes de sa migration. *C. R. Acad. Sci. Paris*, 1949, 228, p. 772.
- FONTAINE, M. - Physiologie du Saumon (suite). *Ann. Stat. Centr. Hydrob. Appl.*, 1950, 3, pp. 7—13.
- , CALLAMAND, O. et VIBERT, R. - La physiologie du Saumon. Deuxième partie. *Ann. Stat. Centr. Hydrob. Appl.*, 1950, 3, pp. 15—26.
- FONTAINE, M. et LELOUP, J. - Sur l'iodémie de deux Téléostéens migrateurs potamotiques *Salmo salar* L. et *Alosa alosa* L. au début de leur montée reproductrice. *C. R. Acad. Sci. Paris*, 1950, 230, pp. 775.
- L'iodémie du Saumon (*Salmo salar* L.) au cours de sa migration reproductrice. *C. R. Acad. Sci. Paris*, 1950, 230, pp. 1216.
- L'iodémie du jeune Saumon (*Salmo salar* L.) en eau douce. *C. R. Acad. Sci. Paris*, 1950, 231, pp. 169—171.
- L'iodémie d'un Cyclostome marin (*Petromyzon marinus* L.) au moment de sa migration reproductrice. *C. R. Acad. Sci. Paris*, 1950, 230, p. 1538.
- FONTAINE, M. - Les facteurs externes et internes dans les migrations des Poissons. Rapport Colloque International d'Ecologie, Paris, février 1950.
- et HATEY, J. - Sur la teneur en glycogène du foie du Saumon (*Salmo salar* L.) adulte, à diverses étapes de sa migration. *C. R. Soc. Biol.*, 1950, 144, p. 788.
- et ———. - Variations de la teneur du foie en glycogène chez le jeune Saumon (*Salmo salar* L.) au cours de la smoltification. *C. R. Soc. Biol.*, 1950, 144, p. 953.
- et RAFFY, A. - Le facteur hypophysaire de rétention d'eau chez les Téléostéens. *C. R. Soc. Biol.*, 1950, 144, p. 6.
- FRANCOIS, Y. - Quelques aspects de la reproduction chez les Poissons. *Bull. Fr. Piscicult.*, 1946, 140, pp. 108—113.

- GABE, M. et VEIL, C. - Sur la présence de basophiles tissulaires dans l'écaille de *Cyprinus carpio*. *J. Physiol.*, 1948, 40, 187A.
- GALLIEN, L. - Sur un cas d'arrhénoïdie chez *Lebistes reticulatus*. *Bull. Soc. Zool. Fr.*, 1945, 70, pp. 104—106.
- Masculinisation des caractères sexuels somatiques et analyse génétique chez *Lebistes reticulatus*. *C. R. Acad. Sci. Paris*, 1946, 222, p. 1522.
- Masculinisation du phénotype sexuel chez *Lebistes reticulatus* et bipotentialité du soma. *Bull. Soc. Zool. Fr.*, 1947, 71, pp. 213—214.
- GALLOIS, C. - L'Alose du Rhône. *Bull. Fr. Piscicult.*, 1946, 141, pp. 162—176.
- L'Alose du Rhône (suite). *Bull. Fr. Piscicult.*, 1946, 142, pp. 5—14.
- L'Alose du Rhône (suite). *Bull. Fr. Piscicult.*, 1946, 143, pp. 72—79.
- L'Alose du Rhône (suite et fin). *Bull. Fr. Piscicult.*, 1947, 144, pp. 130—136.
- La fécondation artificielle de l'Alose du Rhône. *Bull. Fr. Piscicult.*, 1947, 145, p. 198.
- GOSSOT, P. - Ce qu'il faut savoir des Poissons des Eaux douces de la France. Lechevalier éd., Paris, 1946, 257 p., 20 pl. color.
- HOESTLANDT, H. - Fécondation artificielle et incubation chez un Téléostéen: *Paralosa rhodanensis* R. *C. R. Soc. Biol.*, 1947, 141, pp. 937—938.
- Fécondation artificielle et incubation de l'Alose du Rhône: *Paralosa rhodanensis* Roule. *Ann. Stat. Centr. Hydrob. Appl.*, 1948, 2, pp. 223—228.
- HOESTLANDT, H. et VINCENT, H. - Recherches sur l'influence d'une hétéro-auxine (acide 2—4 dichlorophénoxyacétique) chez les Vertébrés Inférieurs. *Ann. Stat. Centr. Hydrob. Appl.*, 1950, 3, pp. 173—184.
- JULLIEN, A., LAURENT, P. et RIPPLINGER, J. - Sur la teneur des principaux organes en acétylcholine chez les Poissons d'eau douce. *Bull. Soc. Hist. Nat. Doubs Besançon*, 1950.
- De l'importance des Mollusques et des Poissons dans les recherches expérimentales sur le cœur et son innervation extrinsèque. *Bull. Soc. Hist. Nat. Doubs Besançon*, 1950.
- KOBIS, G. - Quelques aspects histochimiques du rôle du syncytium vitellin de *Salmo irideus*. Diplôme d'Etudes Supérieures, Fac. Sci. Paris, 21 décembre 1949, 35 p. dactylogr., 2 pl.
- LAFON, M. - Recherches sur l'utilisation des réserves vitellines au cours de l'embryogénèse. Données sur l'embryon de Truite (*Salmo fario* et *Salmo irideus*). *Arch. Intern. Physiol.*, 1947, 55, pp. 123—152.
- LAUR, C. M. - Etude des oxydases dans les leucocytes de quelques Poissons. *Le Sang*, 1947, 18, pp. 170—173.
- Etude des réticulocytes chez certains Poissons et problème de la mégalo blastose. *Le Sang*, 1947, 18, pp. 312—319.
- Etude des leucocytes mononucléaires granuleux chez la Carpe. *Le Sang*, 1949, pp. 452—460.
- LEFEVRE, M. - Contribution à la connaissance de la biologie de la Carpe en étangs clos. *Ann. Stat. Centr. Hydrob. Appl.*, 1948, 2, pp. 51—63.
- LELOUP, J. - Action des antithyroïdiens (aminothiazol et phénylthiourée) sur la cuprémie dans la série des Vertébrés. *Bull. Soc. Chim. Biol.*, 1947, 29, p. 582.
- LHOTTE, R. - Influence des variations de salinité sur le rythme cardiaque de la jeune Anguille (Civelle). *C. R. Soc. Biol.*, 1945, 139, pp. 27—28.
- LIGNIÈRES, R. et HOESTLANDT, H. - L'alimentation artificielle de l'alevin de Truite arc-en-ciel (*Salmo irideus* Gibb.). *Bull. Fr. Piscicult.*, 1949, 154, pp. 18—29.



- L'alimentation artificielle de l'Alevin de Truite Arc-en-Ciel (*Salmo irideus* Gibb.) (suite). *Bull. Fr. Piscicult.*, 1949, 155, pp. 49—65.
- L'alimentation artificielle de l'alevin de Truite Arc-en-Ciel (*Salmo irideus* Gibb.) (suite et fin). *Bull. Fr. Piscicult.*, 1950, 156, pp. 97—109.
- MONOD, TH. - Sur la lecture des écailles de quelques Poissons du Niger moyen. *C. R. Acad. Sci. Paris*, 1945, 220, pp. 629—630.
- Dents de remplacement chez deux *Hydrocion* d'Afrique tropicale. *Conf. Intern. Africanistes de l'Ouest*, 1ère réunion, Dakar, 1945.
- Note sur la lecture des écailles de quelques Poissons du Niger moyen. *Conf. Intern. Africanistes de l'Ouest*, 1ère réunion, Dakar, 1945.
- Sur trois dispositifs de "verrouillage" osseux chez les Poissons. *Conf. Intern. Africanistes de l'Ouest*, 1ère réunion, Dakar, 1945.
- Sur deux Citharinidés du Moyen Niger. *Conf. Intern. Africanistes de l'Ouest*, 2ème réunion, Bissao, 1947.
- Eaux douces tropicales. *Cybiu*, 1947, 1, pp. 5—9.
- Sur l'appareil branchiospinal de quelques Téléostéens. *Rapport XIIIème Congrès Intern. Zool. Paris*, juillet 1948, pp. 330—331.
- Données statistiques et biologiques sur quelques Poissons du Niger Moyen. *Cybiu*, 1949, 4, pp. 4—61.
- Conseils pratiques pour l'étude des eaux douces soudanaises. *Cybiu*, 1949, 4, pp. 73—79.
- Sur l'appareil branchiospinal de quelques Téléostéens tropicaux.
- Notes biologiques sur quelques Poissons du Niger. *Bull. Bull. I. F. A. N.*, 1949, 11, pp. 36—76.
- *Fr. Piscicult.*, 1950, 159, pp. 53—55.
- Notes d'ichtyologie ouest-africaine. *Bull. I. F. A. N.*, 1950, 12, pp. 1—71, 110 fig.
- Sur deux *Garra* d'Afrique Occidentale. *Bull. I. F. A. N.*, 1950, 12, pp. 976—983.
- PALLOT, G., TOURNEUX, G. et TOURNEUX, J. - Les variations sexuelles de l'ilôt pancréatique des Téléostéens. *Bull. Hist. Appl.*, 1946, 23, pp. 112—122.
- PORTAL, J. - Observations sur la pisciculture artificielle du Brochet. *Bull. Fr. Piscicult.*, 1947, 147, pp. 61—70.
- PUYO, J. - Poissons de la Guyane française. Faune de l'Empire Français, 12, O. R. S. O. M., Larose, Paris, 1949, 280 p.
- TAGAND, R. HOESTLANDT, H. et EYRAUD, L. - Etude de quelques facteurs de croissance chez l'alevin de *Salmo irideus* (Gibb.). *Bull. Fr. Piscicult.*, 1947, 145, pp. 172—185.
- THIBAUT, CH. - Action de la lumière sur l'équilibration des Poissons. *C. R. Acad. Sci. Paris*, 1946, 223, pp. 340—342.
- Nouvelles recherches sur le rôle de la lumière dans l'équilibration des Poissons. Eclairissement monochromatique de la rétine centrale de la Carpe. *C. R. Soc. Biol.*, 1947, 141, pp. 378—380.
- Nouvelles recherches sur le rôle de la lumière dans l'équilibration des Poissons. Eclairissements monochromatiques des rétines périphériques chez la Carpe. *C. R. Soc. Biol.*, 1947, 141, pp. 380—382.
- et THIBAUT, O. - Mécanismes physiologiques de l'adaptation chromatique chez les Poissons. *Année Biologique*, 1947, 23, pp. 161—181.
- THIBAUT, CH. - Action de la lumière blanche et monochromatique sur la posture des Poissons Téléostéens. Utilisation de cette action pour l'étude de la vision. *Arch. Sci. Physiol.*, 1949, 3, pp. 101—124.
- THOMAS, L. - Contribution à l'histopathologie comparée des Poissons. Thèse Sci. Paris, 1948, 922 p. + bibliographie très abondante.

- THOMOPOULOS, TH. - L'ébauche thyroïdienne chez la Truite (*Salmo fario* L. et *Salmo irideus* Gibb.). *C. R. Acad. Sci. Paris*, 1948, 227, pp. 1262—1264.
- Morphogénèse de la thyroïde embryonnaire chez la Truite (*Salmo irideus* Gibb. et *Salmo trutta* L.). *Dipl. Etud. Sup., Fac. Sci. Paris*, 1949, 36 p. dactyl., 10 fig.
- Ebauche et morphogénèse embryonnaire de la glande thyroïde chez la Truite (*Salmo irideus* Gibb. et *Salmo fario* L.). *Arch. Anat. Microsc. Morph. Exp.*, 1949, 38, pp. 216—254.
- Sur la thyroïde de *Gymnarchus niloticus* Cuv. (Téléostéen Mormyridé). *Bull. Soc. Zool. Fr.*, 1950, 75, pp. 293—306.
- VIBERT, R. - Les poissons migrateurs dans l'économie piscicole du Sud-Ouest. *Bull. Fr. Piscicult.*, 1945, 136, pp. 121—135.
- Principes d'aménagement d'une rivière à Saumons. *Bull. Fr. Piscicult.*, 1945, 138, pp. 5—23.
- Du repeuplement en Truites et Saumons par enfouissement de boîtes d'alevinage garnies d'oeufs dans les graviers. *Bull. Fr. Piscicult.*, 1949, 153, pp. 125—150.
- Aperçu sur la biologie du Saumon atlantique. *Ann. Stat. Centr. Hydrob. Appl.*, 1948, 2, pp. 145—152.
- Le régression du saumon du Rhin. *Bull. Fr. Piscicult.*, 1950, 156, pp. 131—134.
- Recherches sur le Saumon de l'Adour. *Ann. Stat. Centr. Hydrob. Appl.*, 1950, 3, pp. 27—149.
- VILTER, V. - Métamorphose myo-somatique de larves d'Anguille et relations avec la croissance stomacale. *C. R. Soc. Biol.*, 1945, 139, pp. 808—810.
- Hypophysectomie chez la larve d'Anguille (Civelle) et répercussions sur la métamorphose pigmentaire. *C. R. Soc. Biol.*, 1945, 139, pp. 816—817.
- Inhibition expérimentale de la croissance stomacale au cours de la métamorphose larvaire de l'Anguille. *C. R. Soc. Biol.*, 1945, 139, pp. 817—818.
- Action de la thyroxine sur la métamorphose larvaire de l'Anguille. *C. R. Soc. Biol.*, 1946, 140, pp. 783—785.
- L'évolution pigmentaire des larves d'Anguilles dans ses relations avec la tolérance vis-à-vis de l'eau de mer. *C. R. Soc. Biol.*, 1946, 140, pp. 275—277.
- Déterminisme hypophysaire du comportement halophile de Civelles immigrantes. *C. R. Soc. Biol.*, 1946, 140, pp. 483—484.
- WURTZ-ARLET, J. - Quelques observations sur la biologie de la Vandoise. *Bull. Fr. Piscicult.*, 1950, 158, pp. 14—18.



## BATRACIENS

- ANCEL, P. et VINTEMBERGER, P. - Sur le mouvement de translation en direction dorsale qu'effectue le noyau de segmentation durant la rotation de fécondation, dans l'oeuf de Grenouille rousse. *C. R. Soc. Biol.*, 1947, 141, pp. 553—556.
- Sur la localisation du noyau dans l'hémisphère dorsal de l'oeuf fécondé, chez la Grenouille rousse. *C. R. Soc. Biol.*, 1947, 141, pp. 556—558.
- Recherches sur le déterminisme de la symétrie bilatérale dans l'oeuf des Amphibiens. *Bull. Biol. Fr. Belg.*, 1948, 31, p. 182.
- Sur l'orientation du plan de symétrie bilatérale par la centrifugation de l'oeuf de *Rana fusca*. Oeufs fécondés. *C. R. Soc. Biol.*, 1949, 143, pp. 1313—1314.
- La rotation de symétrisation, facteur de la polarisation dorso-ventrale des ébauches primordiales, dans l'oeuf des Amphibiens. *Arch. Anat. Micr. Morph. Exp.*, 1949, 38, pp. 167—183.
- ANGEL, F. et GUIBÉ, J. - Tableau des espèces actuellement connues du genre *Pseudohemiscus* et description d'une nouvelle espèce: *P. pustulosus*. *Bull. Soc. Zool. Fr.*, 1945, 70, pp. 150—154.
- Etude comparative des espèces *Spinosa*, *Courtoisi* et *Delacouri*, appartenant au genre *Rana*. *Bull. Mus. Nat. Hist. Nat.*, 1945, 17, pp. 379—382.
- ANGEL, F. - Petit Atlas des Amphibiens et des Reptiles. N. Boubée, éd., Paris, 1946.
- Faune de France no 45: Reptiles et Batraciens. Lechevalier, éd. Paris, 1946, 202 p.
- Vie et moeurs des Amphibiens. Payot éd., Paris, 1947, 317 p.
- et LAMOTTE, M. Note sur la biologie d'un Crapaud vivipare, *Nectophrynoïdes occidentalis* Ang. *C. R. Acad. Sci. Paris*, 1947, 224, pp. 413—415.
- † ANGEL, F. - Reptiles et Amphibiens de Madagascar et des îles voisines, faisant partie des collections du Musée Zoologique de Strasbourg. *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, pp. 553—558.
- *Arthroleptis cruscum* et *A. nimbaense*, Batraciens nouveaux de Guinée française (matériaux de la mission LAMOTTE aux Monts Nimba). *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, pp. 559—562.
- ARVY, L., BOIFFARD, J. A. et GABE, M. - Effets des rayons X sur l'hypophyse de *Rana temporaria* L. *C. R. Soc. Biol.*, 1947, 141, pp. 340—341.
- Les modifications de l'hémogramme déterminées chez *Rana temporaria* L. par l'irradiation de l'hypophyse aux rayons X. *C. R. Soc. Biol.*, 1947, 141, pp. 455—456.
- ARVY, L. - Le dimorphisme sexuel sanguin chez *Rana temporaria* L. et *Bufo vulgaris* Laur. *C. R. Soc. Biol.*, 1947, 141, pp. 456—457.
- Action de la thyroxine sur l'hématopoïèse de la Grenouille d'hiver castrée. *C. R. Soc. Biol.*, 1949, 143, pp. 364—365.
- Intervention de la sécrétion thyroïdienne dans le réveil printanier de l'hématopoïèse chez *Rana temporaria* L. *C. R. Acad. Sci. Paris*, 1950, 230, pp. 2118—2120.
- Activation de la glande thyroïde au moment du réveil printanier chez *Rana temporaria* L. *Experientia*, 1950. (sous presse)
- BOGORAZE, D. - Le système nerveux caudal des larves d'Amphibiens Anoures.

- Topographie et architecture du soutien de la moëlle. *C. R. Acad. Sci. Paris*, 1946, 223, pp. 209—210.
- Sur l'existence des racines ventrales, croisées, nées d'un ganglion intramédullaire, et présumées sensibles, chez les têtards d'Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1946, 226, pp. 1217—1219.
- BRUMPT, E. - Utilisation éventuelle du Batracien *Xenopus laevis* dans la lutte contre les larves de Moustiques; interactions diverses à prévoir. *C. R. Acad. Sci. Paris*, 1945, 220, pp. 923—925.
- CAMBAR, R. - Début du fonctionnement du pronéphros chez l'Embryon de Grenouille. *C. R. Soc. Biol.*, 1947, 141, pp. 752—754.
- Valeur fonctionnelle du pronéphros chez le très jeune têtard de Grenouille. *C. R. Soc. Biol.*, 1947, 141, pp. 754—756.
- Détermination de la capacité fonctionnelle précoce du mésonéphros chez le têtard de Grenouille. *C. R. Soc. Biol.*, 1947, 141, pp. 760—761.
- Valeur et modalités du transit de l'eau chez les jeunes larves d'Amphibiens. *C. R. Soc. Biol.*, 1947, 141, pp. 761—763.
- L'ablation précoce du pronéphros n'influence pas la morphogénèse du mésonéphros chez les larves d'Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1947, 225, pp. 592—594.
- Destinée du canal de Wolff après ablation précoce du pronéphros chez les larves d'Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1947, 225, pp. 644—646.
- Démonstration expérimentale du rôle inducteur du canal de Wolff dans la morphogénèse du mésonéphros, chez les Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1947, 225, pp. 823—825.
- Précisions expérimentales sur le mode d'action inductrice du canal de Wolff sur le mésonéphros des Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1947, 225, pp. 895—897.
- Nouvelle preuve expérimentale du rôle inducteur du canal de Wolff sur la morphogénèse du mésonéphros des Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1947, 225, pp. 1019.
- La partie antérieure du canal de Wolff peut, expérimentalement, induire la morphogénèse du mésonéphros chez les Amphibiens Anoures. *C. R. Acad. Sci. Paris*, 1948, 226, pp. 1542—1544.
- Recherches expérimentales sur les facteurs de la morphogénèse du mésonéphros chez les Amphibiens Anoures. *Bull. Biol. Fr. Belg.*, 1948, 82, pp. 214—285.
- Données récentes sur le développement du système pronéphrétique chez les Amphibiens (Anoures en particulier). *Année Biologique*, 1949, 25, pp. 115—130.
- CHAMPY, C. et DEMAY, M. - Le rôle des nerfs et des substances élaborées par eux dans le développement des callosités des Batraciens. *Arch. Zool. Exp. Gén.*, 1949, 86, N. et Rev. no 2, pp. 66—77.
- Gâines périvasculaires et autres adaptations au saut chez les Grenouilles. *C. R. Soc. Biol.*, 1949, 143, pp. 189—190.
- COUTEAUX, R. - Rapports du buisson de Kühne et des noyaux musculaires chez la Grenouille. *C. R. Soc. Biol.*, 1945, 139, pp. 376—377.
- CRÉZÉ, J. - L'invitation à la ponte chez *Xenopus laevis*. *Feuille Nat.*, 1949, 4, pp. 75—76 et 1949, 4, p. 4.
- DELSOL, M. - Utilisation du thiouracile pour évaluer le rôle et l'activité de la thyroïde larvaire chez les têtards de *Discoglossus pictus* Ott. *C. R. Acad. Sci. Paris*, 1950, 230, pp. 237—238.



- GABE, M. - Sur l'autogreffe hépatique chez la Grenouille d'hiver. *C. R. Soc. Biol.*, 1946, 140, pp. 243—244.
- GALLIEN, L. - Caractère ambisexual de la callosité chez *Xenopus laevis* Daudin. *Bull. Soc. Zool. Fr.*, 1947, 72, pp. 192—193.
- Action inhibitrice de la prégnéninolone sur la thyroïde larvaire de *Discoglossus pictus* Ott. *C. R. Soc. Biol.*, 1949, 143, pp. 341—343.
- Dissociation de la chronologie du développement larvaire chez le têtard de *Rana temporaria* L. soumis à l'action de doses élevées d'éthinyl-testostérone. *C. R. Acad. Sci. Paris*, 1949, 229, pp. 1367—1368.
- GALLIEN, L. et MUGARD, H. - Hétéropléidie chez *Pleurodeles walthii* M. *Bull. Soc. Zool. Fr.*, 1950, 75, p. 162 (démonstration).
- GALLIEN, L. - Etude de quelques facteurs intervenant dans l'action des hormones sexuelles au cours de la différenciation du sexe chez les Amphibiens Anoures. *Bull. Biol. Fr. Belg.*, 1950, 84, pp. 341—375.
- GAUTHIER, C. - La naissance du réflexe amoteur chez le jeune Crapaud. *C. R. Acad. Sci. Paris*, 1946, 223, pp. 1026—1027.
- GUIBÉ, J. - A propos du genre *Gephyromantis* (Batracien). *Bull. Mus. Nat. Hist. Nat.*, 1945, 17, pp. 383—384.
- Variations de la ceinture scapulaire chez les espèces appartenant aux genres *Mantipus* et *Plethodontohyla* (Batraciens). *Bull. Mus. Nat. Hist. Nat.*, 1947, 19, pp. 261—264.
- La répartition géographique des Batraciens de Madagascar. *Mém. Inst. Sci. Madagascar*, 1948, 1, pp. 177—179.
- GUIBÉ, J. - Description d'un Batracien nouveau pour la faune malgache, *Platypelis Milloti* n. sp. (Microhylidae). *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, p. 214.
- HATT, P. - Blastotomie, en apparence spontanée, chez le Triton. *Bull. Soc. Zool. Fr.*, 1945, 70, pp. 116—119.
- HOESTLANDT, H. et VINCENT, H. - Recherches sur l'influence d'une hétéro-auxine (acide 2—4 dichlorophénoxyacétique) chez les Vertébrés Inférieurs. *Ann. Stat. Centr. Hydrob. Appl.*, 1950, 3, pp. 173—184.
- JACQUOT, M. - Note préliminaire sur les corrélations biométriques entre la diagnose céphalique et le développement cérébral chez les Batraciens Urodèles, notamment *Triturus helveticus*. *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, pp. 327—332.
- Diagnose céphalique et développement cérébral chez les Urodèles. Comparaison entre *Triturus helveticus* et *Amblystoma tigrinum*. *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, pp. 445—449.
- JOLLY, J. - Sur la fonction du système lymphatique des Batraciens. *C. R. Soc. Biol.*, 1945, 139, pp. 672—674.
- La régénération de la tête chez les Batraciens. *C. R. Soc. Biol.*, 1946, 140, pp. 12—13.
- Recherches sur le système lymphatique des Batraciens. *Arch. Anat. Micr. Morph. Exp.*, 1946—47, 36, pp. 3—44.
- Sur la régénération du globe oculaire chez les Batraciens. *C. R. Soc. Biol.*, 1949, 143, pp. 895—896.
- LALLIER, R. - Action de divers agents oxydants et réducteurs sur les explants dorsaux et ventraux de jeunes gastrulas de Triton. *C. R. Soc. Biol.*, 1949, 143, pp. 736—738.
- LAUR, C. M. - Réaction nouvelle des réticulocytes du sang de Triton. *Le Sang*, 1948, pp. 57—59.

- LECAMP, M. - Recherche du glycogène dans les tissus de régénération chez les Batraciens. *C. R. Acad. Sci.*, 1947, 224, pp. 230—231.
- Influence de substances glucidiques et protéiques sur la régénération des membres chez le Triton. *C. R. Acad. Sci.*, 1947, 224, pp. 363—364.
- et DELSOL, M. - Influence du benzopyrène sur la régénération des ébauches de membres chez le têtard du Crapaud accoucheur. *C. R. Acad. Sci.*, 1947, 224, pp. 499—501.
- LECAMP, M. - Les tissus de régénération en culture in vitro. *C. R. Acad. Sci.*, 1947, 224, pp. 674—675.
- Régénération chez le Triton in vitro et in vivo. *C. R. Acad. Sci. Paris*, 1948, 226, pp. 695—696.
- De l'influence de substances hétérogènes sur le tissu de régénération. *C. R. Acad. Sci. Paris*, 1949, 228, pp. 953—954.
- LHOSTE, J. et ROTH, P. - Sur l'action des solutions aqueuses de 2—4 dichlorophénoxyacétate de sodium sur l'évolution des oeufs de *Rana temporaria* L. *C. R. Soc. Biol.*, 1946, 140, pp. 272—273.
- MILLOT, J. et GUIBÉ, J. - Les Batraciens du Nord de l'Andringitra (Madagascar). *Mém. Inst. Sci. Madagascar*, 1950, A, 4, pp. 197—206.
- MONCEAU, F. - Lieux de terminaison des fibres du pneumogastrique dans le coeur des Batraciens (Grenouille, Crapaud). *C. R. Soc. Biol.*, 1946, 140, pp. 286—288.
- MORON, Y. - Observations sur *Xenopus tropicalis* Gray. *Bull. Soc. Zool. Fr.*, 1947, 72, pp. 128—132.
- PAULIAN, R. et VILARDEBO, A. - Observations sur le régime alimentaire des Batraciens en Basse-Côte d'Ivoire. *Bull. Soc. Zool. Fr.*, 1946, 71, pp. 129—132.
- PICARD, D. et TRUCHMANN-DUPLESSIS, H. - Le système réticulo-endothélial dans la biologie du Triton alpestre. *Bull. Hist. Appl.*, 1946, 23, pp. 93—107.
- RAFFY, A. - Physiologie des animaux amphibiens. *Année Biologique*, 1945, 21, pp. 77—92.
- REY, P. - Sur l'existence de races géographiques chez *Bufo vulgaris* Laur. *Bull. Soc. Zool. Fr.*, 1946, 71, pp. 190—193.
- ROSTAND, J. et MUGARD, H. - Action du gaz carbonique sur les oeufs de Batraciens. *Rev. Sci.*, 1946, 84, p. 292.
- ROSTAND, J. - Essai d'inoculation de noyaux embryonnaires dans des oeufs de Grenouilles irradiées. *Rev. Sci.*, 1946, 84, pp. 412—413.
- Action du froid sur l'oeuf immature de *Rana temporaria*. *Rev. Sci.*, 1947, 85, p. 363.
- De quelques anomalies digitales chez le Crapaud ordinaire (*Bufo bufo*). *Rev. Sci.*, 1947, 85, pp. 1121—1125.
- Conservation à basse température des ovules coelomiques de Grenouille rousse (*Rana temporaria*). *C. R. Soc. Biol.*, 1949, 143, pp. 17—18.
- Inoculation de cellules et de substances chimiques dans l'ovule immature de *Rana*. *C. R. Soc. Biol.*, 1949, 143, pp. 83—84.
- ROTH, P. - Contribution à l'étude de l'action de la thyroxine et des substances antagonistes dans la métamorphose expérimentale des Batraciens Anoures. *Mém. Muséum Hist. Nat.*, 1946, 21, pp. 175—273.
- Sur un cas d'amplexus intergénérique chez un individu mâle de *Rana esculenta*. *Bull. Soc. Zool. Fr.*, 1946, 71, pp. 188—190.
- Sur l'action des hormones sexuelles dans la métamorphose expérimentale de l'Axolotl (*Amblystoma tigrinum* Green) provoquée par la thyroxine (1ère note). *Bull. Mus. Nat. Hist. Nat.*, 1946, 18, pp. 300—305.



- Sur l'action des hormones sexuelles dans la métamorphose expérimentale de l'Axolotl (*Amblystoma tigrinum* Green) provoquée par la thyroxine (2ème note). *Bull. Mus. Nat. Hist. Nat.*, 1947, 19, pp. 131—134.
- Sur l'action antagoniste des substances oestrogènes dans la métamorphose expérimentale des Amphibiens (3ème note). *Bull. Mus. Nat. Hist. Nat. Paris*, 1948, 20, pp. 408—415.
- Sur l'action des hormones sexuelles dans la métamorphose des Amphibiens hypophysoprivés (4ème note). *Bull. Mus. Nat. Hist. Nat. Paris*, 1950, 22, pp. 67—72.
- Etude de l'action de la thiourée, de l'aminothiazol et du méthylthiouracile dans la métamorphose des Batraciens Anoures. *Ann. Endocr.*, 1947, 8, pp. 254—275.
- Sur l'action de l'extrait d'hypophyse, seul ou combiné avec d'autres hormones, sur le développement et la métamorphose de l'Axolotl (5ème note). *Bull. Mus. Nat. Hist. Nat.*, 1950, 22, pp. 657—663.
- SENTEIN, P. - L'oeuf indivis plurinuclée, orientation aberrante du développement provoquée par les antimitotiques. *Bull. Hist. Appl.*, 1948, 25, pp. 193—194.
- Action de diverses substances: caféine, uréthane, cacodylate, trypan-flavine, sur la mitose. *Bull. Hist. Appl.*, 1948, 25, p. 194.
- Production expérimentale d'oeufs insegmentés plurinuclées chez le Triton sous l'influence de substances antimitotiques. *C. R. Soc. Biol.*, 1948, 142, pp. 1426—1428.
- Arrêt de la segmentation, blocage de la mitose et polyploidie par l'action de l'éthyluréthane sur l'oeuf de Batracien. *C. R. Acad. Sci. Paris*, 1949, 228, pp. 706—707.
- SERFATY, A. - La résistance de la Grenouille (*Rana esculenta* L.) à l'asphyxie. *Feuille Nat.*, 1946, 1, pp. 32—33.
- STOLL, R. - Sur quelques phénomènes initiaux du développement de l'oeuf de *Bufo vulgaris*. *C. R. Soc. Biol.*, 1946, 140, pp. 102—103.
- De l'influence du point d'insémination sur la position du plan de symétrie bilatérale de l'oeuf de *Bufo vulgaris*. *C. R. Soc. Biol.*, 1946, 140, pp. 306—307.
- STUTINSKY, F. - Etude de la mélanocinèse chez la Grenouille à l'aide d'une cellule photoélectrique. Action de l'interméline chez la Grenouille normale. *C. R. Acad. Sci. Paris*, 1945, 221, pp. 153—155.
- Température et action de l'interméline sur la mélanocinèse de la Grenouille hypophysectomisée. *C. R. Soc. Biol.*, 1946, 140, pp. 771—773.
- Etude photoélectrique de la mélanocinèse chez la Grenouille en vue de la détection et du dosage de l'interméline. *Congrès de l'Ass. Fr. Av. Sciences*, Nice, sept. 1946.
- Action des injections répétées d'interméline sur la mélanocinèse de la Grenouille. *C. R. Soc. Biol.*, 1947, 141, pp. 1034—1036.
- THOMAS, J. A., ROSTAND, J., GRÉGOIRE, J. - Action inhibitrice de doses minimales de ribonucléase sur la segmentation de l'oeuf de la Grenouille rousse. *C. R. Acad. Sci. Paris*, 1946, 222, pp. 1139—1141.
- THOMAS, J. A. et BORDERIOUX, I. - La survie expérimentale d'Urodèles en milieu synthétique, antiseptique et nutritif. *C. R. Acad. Sci. Paris*, 1947, 225 pp. 963—964.
- Recherches sur la survie expérimentale d'organites chez les Urodèles en milieu synthétique, antiseptique et nutritif. *Arch. Anat. Microsc. Morph. Exp.*, 1948, 37, pp. 263—288.

- TUCHMANN-DUPLESSIS, H. - Variations saisonnières et répercussions de l'hypophysectomie sur la thyroïde du Triton alpestre. *Bull. Hist. Appl.*, 1945, 22, pp. 17—25.
- Régénération d'une patte surnuméraire chez un Triton après amputation de la patte du côté opposé. *Bull. Soc. Zool. Fr.*, 1945, 70, pp. 159—161.
- Régénération chez le Triton. Influence de l'hypophysectomie et de l'introduction parentérale de substances médicamenteuses. *Bull. Hist. Appl.*, 1946, 23, pp. 171—183.
- Action des iodoprotéines sur la thyroïde et la métamorphose des Batraciens. *Bull. Hist. Appl.*, 1948, 25, pp. 132—136.
- Action de l'hormone gonadotrope et lactogène sur le comportement et les caractères sexuels secondaires du Triton normal et castré. *Arch. Anat. Microsc. Morph. Exp.*, 1949, 38, pp. 302—317.
- Action des sels de béryllium sur la régénération chez le Triton. *Bull. Hist. Appl.*, 1950, 27, pp. 116—117.
- TUSQUES, J. - Action de la thyroxine sur l'appareil olfactif des têtards de *Rana esculenta*. *C. R. Soc. Biol.*, 1949, 143, pp. 245—247.
- Action de la thyroxine sur l'appareil visuel des têtards de *Rana esculenta*. *C. R. Soc. Biol.*, 1949, 143, pp. 332—334.
- Corrélation entre le développement des membres et le développement des structures nerveuses centrales correspondantes sous l'action de la thyroxine chez les têtards de *Rana esculenta*. *C. R. Soc. Biol.*, 1949, 143, pp. 380—382.
- Insensibilité à la thyroxine du régénérat de nageoire caudale des têtards d'Anoures. *C. R. Soc. Biol.*, 1949, 143, pp. 648—650.
- VALLÉE, L. - Contribution à l'étude des races sexuelles françaises de *Rana temporaria* L. *Bull. Soc. Zool. Fr.*, 1947, 72, pp. 97—100.

Adresse de l'auteur:  
LABORATOIRE DES PÊCHES COLONIALES  
Gif-sur-Yvette, Seine et Oise

# Taxonomy of the nymphs of the British species of *Leptophlebiidae* (Ephem.)

T. T. MACAN

(Freshwater Biological Association, Ferry House, Ambleside,  
Westmorland, England).

(with 4 figures)

As in previous studies (see Macan 1951 for references), specific distinction has been established on cast skins identified from the adults that emerged from them. Points of difference having been found in this way, it has been possible to base the descriptions on both cast skins and whole nymphs.

There are six British species in this family: two in the genus *Leptophlebia*, three in the genus *Paraleptophlebia*, and one in the genus *Habrophlebia*.

The nymphs of all have been described before, but critical comparison of the species has revealed differences not noticed previously and shown that some of the distinctions used hitherto are less reliable than has been thought.

## *Separation of the genera*

*Habrophlebia* is distinguished at once by its gills, each of which has several filaments (fig. 3F), and there are other points of difference, which have been mentioned in the description of *H. fusca*. *Leptophlebia* and *Paraleptophlebia* are also distinguished by the gills, though otherwise they resemble each other closely. The gills of *Leptophlebia* are swollen at the base (fig. 3M, V), those of *Paraleptophlebia* are more or less straight-sided (fig. 3S, T). In small specimens of *Leptophlebia*, however, the basal swelling of the gill is very slight and in specimens 3 mm. long (the smallest I have seen)






Fig. 1

V, F maxilla of *Leptophlebia vespertina* and *Habrophlebia fusca*. V labrum of *L. vespertina*. S, C maxillary palp and outline of maxilla of *Paraleptophlebia submarginata* and *P. cincta*. f, v labium of *H. fusca* and *L. vespertina*.

it is easily overlooked unless the gill is examined carefully. The maxillary palps of *Paraleptophlebia* (fig. 1S, C) are longer than those of *Leptophlebia* (fig. 1V). Specimens of *Paraleptophlebia submarginata* 3 mm. long (small specimens of the other two species in this genus were not available) are more easily distinguished from specimens of *Leptophlebia* of the same size by the maxillary palps than by the gills.

### *Descriptions and comparisons of the species*

#### *Leptophlebia vespertina* (Linnaeus)

*Previously described by* Lestage 1917 (p. 330) and 1919 (p. 120 and fig. 7).

*Material:* Blelham Beck, WL, 2 cast skins; Cunsey Beck, WL, 1 cast skin; Nor Moss Beck, WL, 6 cast skins; Waterston Moss, WL, 4 cast skins; Windermere, WL, 1 cast skin; Wise Een Tarn, WL, 2 cast skins; Wray Mires Tarn, WL, 3 cast skins; Buttermere, CU, 2 cast skins; Wicken Fen, CB, 6 whole nymphs; Ditchend Brook, SH, 6 whole nymphs; Dockens Water, SH, 1 cast skin.

*Markings:* The nymph is brown. On each abdominal tergite there is a pair of triangular light dots near the proximal margin, and larger less distinct light areas near the sides. This pattern is rather constant, though the degree of contrast between the light and dark parts varies considerably.

*Lengths:* 7.0—10.0 mm.

*Mouthparts, legs, and gills* are illustrated in figs. 1, 2, 3 and 4. It will be more convenient to discuss them under the next species.

*Ecology:* In the Lake District (WL and CU) this species abounds in the more sheltered parts of lakes (Moon 1936) and in tarns (Macan 1949) and occurs also in slowly flowing streams where there is vegetation. In southern Hampshire (SH), nymphs were found in streams draining the sandy base-deficient soils of the New Forest, but not in the calcareous waters of the River Avon and its tributaries (Macan, T.T. and Z. 1939). The species was found in ditches among the sedge at Wicken Fen, a region of calcareous peat.

*Leptophlebia marginata* (Linnaeus).

*Previously described* by Lestage (1917, p. 328 and fig. 25).

*Material:* Nor Moss Beck, WL, 1 cast skin; Rather Heath 6, WL, 3 cast skins; Three Dubs Tarn, WL, 1 cast skin; Cooper's Bottom, SH, 4 whole nymphs; Loch of Lauriston, 2 whole nymphs. (The last place is presumably in Scotland, but there is no other data on the label and I cannot remember who was kind enough to send these specimens).

*Markings:* The pattern on the abdominal tergites is basically the same as that of *L. vespertina*, but more variable. The lateral pale marks are commonly smaller, and there is often a distal median light dot, line, or triangular mark.

*Lengths:* 10.5—11.5 mm.

*The mouthparts* are identical with those of *L. vespertina* apart from some small details of trichiation: the hairs on the front margin of the




Fig. 2.

Mandibles of *Habrophlebia fusca* (F) and *Leptophlebia vespertina* (V).



labrum and at the tip of the labial palp are somewhat longer, and more of the paraglossae is covered with hairs.

*Gills:* The blade of each gill, except the first, is broad and ovate, sweeping round to meet the filament somewhat abruptly (fig. 3M), whereas that of *L. vespertina* tapers gradually into the filament (fig. 3V). The filament of *L. marginata* is shorter than that of *L. vespertina*. This difference in gill-shape has hitherto been deemed sufficient to distinguish the species (eg. Schoenemund 1930, Ulmer 1943). The shape of the gill, however, depends on its size, and the typical ovate shape of *L. marginata* is found only on full-grown nymphs; moreover, the length of the filament is variable. A specimen of *L. marginata* 8 mm. long has gills very like those of *L. vespertina* of the same size, and the two cannot be separated on this character.

*Legs:* The underside of the fore femora of *L. vespertina* (fig. 4V) is beset with spines most of which are complex and of the type shown in fig. 4v. They are fairly obvious under the low power of the compound microscope (x 165). Some of them have more lateral branches than those illustrated, and a few have numerous tiny lateral branches that are no more than hairs.

The corresponding spines of *L. marginata* appear simple under low power magnification, but under high power (x 750) are seen to be fringed with tiny hairs (fig. 4m). This is the most certain distinction between the species.

On a specimen of *L. vespertina* 3 mm. long, most of the spines had more lateral branches than those of bigger specimens, and there were more spines bordered with hairs, but it was just possible to make out with low power magnification that some of the spines were complex. The smallest specimen of *L. marginata* seen was 5.2 mm. long. The lateral hairs on the spines were rather more conspicuous than those of larger specimens, but they were still detectable only under high power magnification.

The middle legs of the two species do not differ significantly. The upper surface of the tibia of *L. vespertina* usually has a few spines on it, that of *L. marginata* is usually bare except for some long hairs. The tibiae of the hind legs of the two are similar, with fairly numerous spines along the upper surface, but the tarsi differ. Those of *L. marginata* usually have a number of spines along the upper surface (fig. 4M), those of *L. vespertina* none or only a few. Unfortunately this is not invariable, and *L. vespertina* may bear many spines.

The teeth approach nearer the tip of the claw in *L. vespertina* (fig. 4V) than in *L. marginata* (fig. 4M). This difference is constant on the specimens that have been examined and, with a little practice, affords a satisfactory means of distinguishing the two species.

*Ecology:* This species has generally been found with *L. vespertina*, though in smaller numbers.

*Paraleptophlebia submarginata* (Stephens)

*Material:* Nor Moss Beck, WL, 3 cast skins and 2 whole nymphs; R. Rawthey, MY, 4 cast skins and 1 whole nymph; R. Ribble, MY, 1 cast skin; Teal Burn, KF, 3 whole nymphs; Daggons Brook, SH, 2 whole nymphs.

I am indebted to Mr D. Scott of the Scottish Brown Trout Research Laboratory for the specimens from Teal Burn.

*Markings:* Some specimens have abdominal markings similar to those of *L. vespertina*; others are uniformly brown. Some have a pale transverse line and a U-shaped mark at the distal end of the segments.

*Lengths:* 8.0—11.5 mm.

*The mouthparts* are like those of *L. vespertina*, except that the last segment of both sets of palps bears long fine hairs, those on the maxillary palp being particularly long and dense. The whole palp is of unusual length (fig. 1S).

*Gills:* There is a progressive but slight diminution in the size of the gills from the middle to the ends of the series, except that the first is distinctly smaller than the rest, usually about half the size (fig. 3S).

*Legs:* Spines are absent from the upper surface of all the tarsi and from the fore and mid-tibiae.

The spines on the femora occur in three irregular rows, along the top, along the middle, and along the bottom (the leg being in the position shown in fig. 4). On the fore femora most of the spines are fairly long, those in the middle are finer than the others, and all are tapering and pointed except a few at the distal end of the upper row. These taper but are blunt at the tip. The mid femur is similar but there are more blunt spines, most of those in the middle row being blunt. The hind femur has characteristic spines. All those of the middle and bottom rows are blunt and parallel-sided, and some are short (fig. 4s).

The teeth extend a little more than half-way along the claws.



Fig. 3.

M, V 4th gills of *Leptophlebia marginata* and *vespertina*. S, T 1st and 2nd gills of *Paraleptophlebia submarginata* and *tumida*. F 4th gill of *Habrophlebia fusca*.



*Ecology:* This species occurs in sluggish streams and occasionally in rapid stony ones and in rivers.

*Paraleptophlebia tumida* Bengtsson

*Previously described* and figured by Ulmer (1943, p. 336, figs. 1—10).

*Material:* Allen R., SH, 1 cast skin and 11 whole nymphs; R. Till, SW, 1 cast skin and 7 whole nymphs.

*Lengths:* 8—9 mm.

*Distinction from P. submarginata:* 1. The gills diminish slightly and regularly in size from the middle outwards, and the first gill is not distinctly smaller than the rest (fig. 3T).

2. The maxillary palp is not as long, and the hairs on its last segment are fewer and shorter (fig. 1C).

3. All the spines on the hind femora are pointed, and none is very short (fig. 4t).

4. There are fewer fine hairs on the palps of the labium, and the glossae are more pointed.

*Ecology:* Allen River and River Till are streams in chalk-down valleys. They flow with moderate speed over stony beds and are densely overgrown with vegetation. Surface water probably disappears every summer. In these two streams, *P. tumida* was abundant and no other Ephemeroptera were found (T.T. and Z. Macan 1940). The species has never been recorded anywhere else in Britain.

(There is an error in the paper referred to. Allen River was dry on 6:vi, not 6:v as shown in table 1 on p. 59).

*Paraleptophlebia cincta* (Retzius)

*Previously described* and figured by Eaton (1888, pl. 32) and Lestage (1919, p. 342 and fig. 27).

*Material:* R. Wharfe, MY, 1 cast skin; R. Kent, WL, 1 whole nymph.

*Length:* 7.5 mm.

*Distinction from foregoing species:* This species resembles *P. tumida* closely, and both differ from *P. submarginata* in the same way,

though, as will be seen presently, the femoral spines of the two are not identical.

*P. cincta* and *P. tumida* can be separated on the following characters:

1. The spines of the middle and lower rows on the hind femora of *P. cincta* are intermediate in character between those of the other two species; they taper like those of *P. tumida*, but are blunt-ended like those of *P. submarginata* (fig. 4c).

2. The teeth do not extend far beyond the middle of the claw of *P. cincta* (fig. 4C), those of *P. tumida* reach nearly three-quarters of the way to the tip (fig. 4T). The number of teeth is unexpectedly variable and not always higher on *P. tumida*, as might be expected. Six specimens of *P. tumida* had respectively 22, 22, 27, 27, 27, and 28 teeth on one of the hind claws; both specimens of *P. cincta* had 21.

3. The maxillary palp of *P. cincta* is slightly longer.

*Ecology*: This species appears to be fairly widespread but nowhere abundant. In addition to my own two records, I have nine kindly supplied by Mr D. E. Kimmins; all are from rivers.

### *Habrophlebia fusca* (Curtis)

*Previously described* and figured by Eaton (1888, pl. 36) and Lestage (1919, p. 348 and fig. 30).

*Material*: Hog House Beck, WL, 1 cast skin; R. Ribble, MY, 3 cast skins; Daggons Brook, SH, 4 whole nymphs.

*Markings*: The abdominal tergites are uniformly pigmented.

*Lengths*: 6.5—8.0 mm.

*The mouthparts* are of the same type as those of the other two genera, but differ in detail: the front margin of the labrum is more deeply indented; the mandibles are more rounded along the outside (fig. 2F); the maxilla is shorter, and the short conical last segment of its palp is characteristic (fig. 1F); the labial palps are shorter (fig. 1f).

*Gills*: Each is divided into two lobes which bear between them some 5 to 12 filaments according to their position in the series, those in the middle being the most richly provided. No other British Ephemeropteran nymph has gills like this.




Fig. 4.

**M** hind-leg of *Leptophlebia marginata* showing spines on tibia and tarsus; **F** fore-leg of *Habrophlebia fusca*; **V** fore-femur of *Leptophlebia vespertina*. m, v, spines from underside of fore femur of *L. marginata* and *vespertina*; s, c, t spines from underside of hind femur of *Paraleptophlebia submarginata*, *cincta*, *tumida*. T, C, V, M hind claws of *P. tumida*, *P. cincta*, *L. vespertina*, *L. marginata*.



*The legs* are less thickly beset with spines than those of the other species and spines are absent from the upper surface of all tibiae and tarsi (fig. 4F). There are about 12 teeth on the hind claws.

*Ecology*: Hog House Beck, in the Lake District, is a slow stream flowing over a sandy bottom with a few stones and some tufts of vegetation. *Habrophlebia* nymphs are quite common on the stones but none have been taken by sweeping through the weeds with a net. Miss Hampshire found numbers of nymphs on stones in a similar stream in South Lancashire (SL); she did not collect in the vegetation. Daggon's Brook, in the south of England, is also a small slow stream with a sandy bottom in which occasional patches of vegetation are rooted. There are no stones but packets of dead leaves occur at intervals. Nymphs were abundant in these packets of dead leaves but not in the vegetation. *H. fusca* has been taken in various other streams and some rivers, but these were the only places where it was numerous.

CB = Cambridgeshire, CU = Cumberland, MY = Mid Yorkshire, SH = South Hampshire, WL = Westmorland (England), KF = Kinross and Fifeshire (Scotland).

*Acknowledgments*

It is a pleasure to acknowledge my indebtedness to Mr D. E. Kimmins who has identified many of the adults and put the records of the British Museum freely at my disposal; to my assistant, Miss Jean Hampshire, who has drawn some of the figures and helped with all aspects of the work; and to Dr J. Senez who has translated the key into French.

*Key*

- 1. Gills with several branches (fig. 3F) . . . . *Habrophlebia fusca*  
Gills with two branches (fig. 3M, V, S, T) . . . . . 2
- 2. Gills (except the first) expanded at the base (fig. 3M, V)  
*Leptophlebia* 3
- Gills straight-sided (fig. 3S, T) . . . . . *Paraleptophlebia* 4
- 3. Compound spines on the underside of the fore femora (fig. 4v); teeth extending about five-sixths of the way

- along the claw (fig. 4V); blade of gills tapering and merging gently into the filament (fig. 3V) ..... *L. vespertina*
- Finely frayed spines, appearing simple except under high power magnification, on the underside of the fore femora (fig. 4m); teeth extending about three-quarters of the way along the claw (fig. 4M); blade of gills strongly rounded, meeting the filament abruptly (fig. 3M) (only in full-grown specimens) ..... *L. marginata*
  - 4. First gill distinctly smaller than all the rest (fig. 3S); spines on underside of hind femora parallel-sided and blunt at the tip, many of them short (fig. 4s); maxillary palps large, the last segment densely covered with long fine hairs (fig. 1S) ..... *P. submarginata*
  - First gill only slightly, if at all, smaller than the others (fig. 3T); spines on underside of hind femora tapering (fig. 4c, t); maxillary palps smaller, the last segment less densely covered with long fine hairs (fig. 1C) ..... 5
  - 5. Spines on underside of hind femora pointed, rather few of them short (fig. 4t); teeth extending nearly three-quarters of the way along the claw (fig. 4T) ..... *P. tumida*
  - Spines on underside of hind femora tapering but blunt at the tip (fig. 4c); teeth not extending far beyond the middle of the claw (fig. 4C) ..... *P. cincta*

#### *Tableau synoptique*

1. Branchies multi-ramifiées (fig. 3F) ..... *Habrophlebia fusca*
- Branchies bifides (fig. 3M, V, S, T) ..... 2
2. Branchies renflées à la base (sauf la première) (fig. 3M, V)  
*Leptophlebia* 3
- Branchies à bords rectilignes (fig. 3S, T) *Paraleptophlebia* 4
3. Epines très dentelées (fig. 4v) sur la face inférieure des fémurs antérieurs; la rangée de dents s'étend sur les 5/6 environ de la griffe (fig. 4V); le renflement basal de la

- branchie se rétrécit progressivement vers l'extrémité (fig. 3V) ..... *L. vespertina*
- Epines simples (avec fines dentelures seulement visible aux forts grossissements) (fig. 4m) sur la face inférieure des fémurs antérieurs; la rangée de dents s'étend sur les 3/4 environ de la griffe (fig. 4M); le renflement basal des branchies est de forme ovale et se termine brusquement (fig. 3M) (ce caractère n'existe que sur les nymphes ayant atteint leur complet développement) ..... *L. marginata*
  - 4. Première branchie beaucoup plus courte que les autres (fig. 3S); sur la face inférieure des fémurs postérieurs, épines à bords rectilignes et à terminaison brusque, souvent courtes (fig. 4s); palpes maxillaires longues, recouvertes, sur leur dernier segment, par un grand nombre de poils longs et fins (fig. 1S) ..... *P. submarginata*
  - Première branchie de même longueur, ou à peine plus courte, que les autres (fig. 3T); sur la face inférieure des fémurs postérieurs épines effilées (fig. 4c, t); palpes maxillaires plus courtes, recouvertes, sur leur dernier segment, par un moins grand nombre de poils longs et fins (fig. 1C) 5
  - 5. Sur la face inférieure des fémurs postérieurs, épines effilées, à extrémités pointues, rarement courtes (fig. 4t); la rangée de dents atteint presque les 3/4 de la griffe (fig. 4T) ..... *P. tumida*
  - Sur la face inférieure des fémurs postérieurs, épines effilées mais à extrémités tronqués (fig. 4c); la rangée de dents dépasse à peine la moitié de la griffe (fig. 4c) ..... *P. cincta*

#### REFERENCES

- EATON, A. E. - 1883—1888. - A revisional monograph of recent Ephemeridae or mayflies. *Trans. Linn. Soc. Lond. Zool.* 3, 1—352.
- LESTAGE, J. A. - 1917, 1919. - Contribution à l'étude des larves des Ephémères paléarctiques (ser. 1 and 2). *Ann. Biol. lacust.* 8, 213—458; 9, 79—182.
- MACAN, T. T. - 1949. - Survey of a moorland fishpond. *J. Anim. Ecol.* 18, 160—186.
- 1951. - The taxonomy of the British species of Siphonuridae. *Hydrobiol.* 3, 84—92.



- and Z. - 1940. - Preliminary note on the Ephemeroptera and Plecoptera of the Hampshire Avon and its tributaries. *J. Soc. Brit. Ent.* 2, 53—61.
- MOON, H. P. - 1936. - The shallow littoral region of a bay at the north-west end of Windermere. *Proc. Zool. Soc. Lond.* 491—515.
- SCHOENEMUND, E. - 1930. - Eintagsfliegen oder Ephemeroptera in: *Die Tierwelt Deutschlands*. 19, 1—106.
- ULMER, G. - 1943. - Die von Prof. A. Thienemann in der Umgegend von Abisko (Lappland) gesammelten Eintagsfliegen und ihre Larven. *Arch. Hydrobiol.* 40, 329—361.

Address of the author:  
FRESHWATER BIOLOGICAL ASSOCIATION,  
Ferry House, Ambleside,  
West-Morland, England.

The algal flora of southeastern  
United States VI.  
Additions to our knowledge of the  
desmid genus *Euastrum* 2.

ARTHUR M. SCOTT

New Orleans Academy of Sciences.

and

GERALD W. PRESCOTT

Michigan State College

Herein are presented two new species and a number of new varieties and forms of existing species, which have been found in gatherings made since the publication of our previous paper on *Euastrum* (1945). Nearly all of them, as will be noted, come from the State of Florida, the rich desmid-flora at which has previously been commented upon by us and other authors. The number of *Euastra* known from southeastern United States now totals 56 species, 59 varieties, and 18 named forms.

We wish to express our thanks to Dr. Hannah Croasdale for translating the diagnoses into Latin, and to Mrs. Dorothy Perine, Jr., for inking the senior author's pencil drawings.

All measurements are given in microns, and the following abbreviations are used: L. = length; W. = width; T. = thickness; I. = isthmus. The symbols (Fla. 25) refer to the senior author's collections, duplicates of which have been deposited in the Farlow Herbarium, Harvard University.

*Euastrum acmon* West & West var. TURGIDUM var. nov. Pl. I, Fig. 10. A variety smaller than the typical and with two prominent protrusions on the median facial protuberance; polar incision broader and more concave than in the typical; in lateral view semicells

transversely elliptic, the lateral margins greatly distended so that the cell is approximately as thick as the semicell is high; in vertical view broadly oval to almost circular in outline, the poles narrowly rounded and with two prominent marginal protrusions on each side in the midregion. L. 30—31; W. 22; T. 18—19; I. 5—6.

Varietas minor quam planta typica, duos protrusiones prominentes in protuberatione media faciali habens; incisio polaris latior et magis concava quam in planta typica; semicellulae a latere visae transverse ellipticae, marginibus lateralibus multum distensis, ita ut cellula quasi aequae crassa ac semicellula alta sit; semicellulae a vertice visae late ovatae ad fere circulares, polis anguste rotundatis, duos protrusionibus marginalibus prominentibus utrimque media in parte visis. Long. 30—31; lat. 22; crass. 18—19; isthm. 5—6.

The specific form and var. *clausum* West and West were originally found in Angola, West Africa, and apparently have not been recorded since.

Florida: Cypress swamp 12.5 miles north of Port St. Joe, Gulf Co. (Fla. 25). Pond 20 miles east of Panama City, Bay Co. (Fla. 47).

*Euastrumbidentatum* var. *oculatum* (Istv.) Krieg. fa. GLABRUM fa. nov. Pl. III, Fig. 1.

A form differing from the variety by having the supraisthmian protuberance smooth and evenly rounded, and by having more or less prominent depressions in the face of the semicell just within the margin of each lateral invagination of the wall; in lateral view semicell inversely cone-shaped from a widely open median incision of the cell, with a smoothly rounded protuberance near the base of the semicell and a slight bulge about half-way between the basal protuberance and the somewhat narrowly rounded apex; a mucilage pore showing between the basal protuberance and the intermediate protuberance. L. 60—61; W. 36—37; T. 25; I. 9—10.

Forma a planta typica differens protuberatione supraisthmali levi atque aequaliter rotundata, et possessione depressionum plus minusve prominentium in superficie semicellulae admodum intra marginem cuiusque invaginationis lateralis membranae; semicellula a latere visa inverse conica ab incisione media late aperta cellulae; et praebens protuberationem leviter rotundatam ad basim semicellulae necnon gibbum circa media in parte inter protuberationem basalem atque apicem quasi anguste rotundatam; porus mucosus inter protuberationem basalem atque protuberationem intermediam visus. Long. 60—61; lat. 36—37; crass. 25; isthm. 9—10.

Florida: Ditch 6 miles south of Punta Gorda, Charlotte Co. (Fla. 76). Swamp 7 miles west of Melbourne, Brevard Co. (Fla. 131).



*Euastrum bidentatum* Naeg. var. **QUADRIOCULATUM** var. nov. Pl. I, Fig. 4.

A variety differing from the typical by having two pores within the lateral margins on each side of the semicell in the mid-region, in addition to the two pores in the face of the semicell; and also differing by having a reduction in the number and size of granules on the wall; central facial protuberance reduced to a low swelling; in lateral view showing six mucilage pores, one on either side in the mid-region and four pores quadrately placed around the center of the ovate semicell; in polar view quadrangularly oval, the lateral margins parallel or slightly convex and with a lateral median protuberance which is without decorations, the poles abruptly rounded, four mucilage pores showing, two on either side just within the margin of the central protuberance. L. 56; W. 36; T. 22; I. 8.

Varietas a planta typica differens accessione duorum pororum intra margines laterales utroque in latere semicellulae media in parte, ad duos poros in superficie semicellulae; differens necnon reductione numeri magnitudinisque granulorum in membrana; protuberatio centralis facialis in inflationem humilem reducta; semicellula a latere visa praebens 6 poros mucosos, unum utrimque media in parte, necnon 4 circum centrum semicellulae ovatae quadrate positos; semicellula a polo visa quadrangulariter ovata, marginibus lateralibus parallelis vel paululum convexis et protuberantem mediam lateralem sine decorationibus praebens; polis abrupte rotundatis, quattuor poris mucosis apparentibus; duo utrimque admodum intra marginem protuberantem centralis. Long. 56; lat. 36; crass. 22; isthm. 8.

Florida: Ditch 10 miles southeast of Port St. Joe, Gulf Co. (Fla. 59).

This variety should be compared with *E. oculatum* Boerges. which it resembles in shape of cell as seen in side view. Like that species also the Florida plants have almost smooth walls. But the form of the basal angles and the deep polar incision identify this plant with *E. bidentatum*.

*Euastrum Ciastonii* Racib. var. **APERTISINUATUM** var. nov. Pl. I, Fig. 5. A variety differing from the typical by having a sinus that opens outwardly from a sharply rounded apex; also differing in its possession of additional granules just within the polar margins of the apical lobules. L. 40—44; W. 27; I. 6.

Varietas a planta typica differens possessione sinus ab apice acute rotundato extrorsum aperti; necnon differens possessione granulorum additiorum admodum intra margines polares apicalium. Long. 40—44; lat. 27; isthm. 6.

Florida: Small lake 12 miles north of Lynn Haven, Bay Co. (Fla. 35).

*Euastrum Ciastonii* var. **ASYMMETRICUM** var. nov. Pl. I, Fig. 6. A variety differing from the typical by having the lower lateral margins of the basal lobes distinctly diverging to rather abruptly rounded lateral angles, the upper lateral margins more deeply retuse, and in having a crest of three granules just within the polar margin of the apical lobules on either side of the median notch; in the face of the semicell differing by having two protuberances in the mid-region which bear two to four granules, one protuberance being distinctly larger than the other; oval in end view, the poles with a marginal series of four granules, with two protuberances in the mid-region on each side, one of which is larger than the other and bidentate rather than simple, and in position diagonally opposite the bidentate protuberance on the other side of the cell. L. 45; W. 29—30; T. 21; I. 8.

Varietas a planta typica differens marginibus lateralibus inferioribus loborum basaliū ad angulos laterales quasi abrupte rotundatos distincte divergentibus, marginibus lateralibus superioribus profundius retusis; differens necnon possessione cristae 3 granulorum admodum intra marginem polarem lobulorum apicalium utroque in laterale incisurae mediae; differens necnon media in facie semicellulae possessione 2 protuberationum 2—4 granula ferentium, una protuberatione admodum maiore quam altera; semicellula a polo visa ovata, polis series marginales 4 granulorum habentibus, et 2 protuberationibus utrimque media in parte, una maiore quam altera et bidentata, non simplici, et positu protuberationi bidentatae altero in latere semicellulae diagonaliter opposita. Long. 45; lat. 29—30; crass. 21; isthm. 8.

Mississippi: Swamp 2 miles west of Pascagoula, Jackson Co. (Miss. 88).

*Euastrum crassum* (Bréb.) Kuetz. Pl. II, Fig. 1.

In southeastern United States the specific form of this plant is much rarer than the several varieties. The form figured herewith has the proportions of var. *microcephalum* Krieg. and var. *michiganense* Presc., but the poreless wall and the facial protuberances of the typical plant. The wall instead of being coarsely scrobiculate is rather finely punctate. L. 179; W. 77; T. 57; I. 24.

Florida: Swamp 3 miles east of Bruce, Walton Co. (Fla. 11).

*Euastrum crassum* var. *scrobiculatum* Lund. fa. Pl. II, Fig. 3.

This form has the lateral margins of var. *microcephalum* Krieg., but the apical lobe and depth of sinus between the polar lobe and the basal lobe which are characteristic of var. *scrobiculatum*. Like

the latter it possesses a central mucilage pore in the face of the semicell. L. 157; W. 76; I. 25.

Florida: Pond 6 miles south of Masaryk, Hernando Co. (Fla. 110).

*Euastrum denticulatum* var. *rectangulare* West & West, fa. Pl. III, Fig. 9.

A variety in which the poles of the cell are approximately as wide as (or a little wider than) the base of the semicell. The granules about the median notch are lacking in the Florida specimens and the lateral margins are more deeply invaginated below the apex than described for the typical. L. 19; W. 15; T. 11; I. 5.

Florida: Swamp 14 miles west of Torrey, Hardee Co. (Fla. 114).

*Euastrum evolutum* var. *integrius* West & West fa. TURGIDUM fa. nov. Pl. III, Figs. 4, 5.

A form in which there is a reduction in the teeth along the margin of the basal lobes and which in lateral view shows the upper lateral margins broadly convex and sloping to a rather sharply rounded apex, with a low protuberance on each side just below the pole and with the median protuberance less prominently developed than in the typical. L. 50—53; W. 36—39; T. 21; I. 8—9.

Forma nota dentium reductione secum dum marginem loborum basalium et, a latere visa, marginibus lateralibus superioribus late convexis et ad apicem quasi acute rotundatum devectis, protuberatione depressa utrimque admodum infra polum, protuberatione media minus prominenter effecta quam in varietate typica. Long. 50—53; lat. 36—39; crass. 21; isthm. 8—9.

Florida: Ditch 4 miles east of LaBelle, Hendry Co. (Fla. 129).

Louisiana: Swamp 4.5 miles south of Ponchatoula, Tangipahoa Parish (La. 111).

The shape of the polar lobe (with subparallel margins) seems to identify this plant with var. *integrius*, but in lateral view it is relatively thicker and is convex along the lateral margins where that variety is retuse. Pl. III, Fig. 5 shows a form from Louisiana approaching var. *trilobum* (Boerges.) Krieg. in the shape of the base of the semicell, but has the apical lobe of var. *integrius*. The former variety is without mucilage pores.

*Euastrum evolutum* var. *monticulosum* (Tayl.) Krieg. fa. PORIFERUM fa. nov. Pl. III, Fig. 2.

A form differing from the typical by having two large mucilage pores in a transverse line just above the facial protuberance which is triangular, and in having the wall coarsely scrobiculate in the mid-



region of the semicell, especially just below and around the facial protuberance. L. 71—78; W. 48—51; T. 28—34; I. 12.

Forma a planta typica differens possessione duorum pororum mucosorum magnorum linea in transversa admodum supra protuberationem facialem, quae triangularis est; differens necnon membrana media in parte semicellulae grosse scrobiculata, praecipue admodum infra circaque protuberationem facialem. Long. 71—78; lat. 48—51; crass. 28—34; isthm. 12.

Florida: Ditch 2.5 miles southeast of Myakka City, Manatee Co. (Fla. 115).

*Euastrum evolutum* (Nordst.) West & West var. REDUCTUM var. nov. Pl. III, Fig. 3.

A variety differing from the typical by having the lateral lobules of the semicell much reduced and with shallow invaginations of the lateral margins rather than incisions; the spinescence of the polar lobules also reduced; in lateral view pole truncate with a protuberance on each side but broadly convex at the apex. L. 58—60; W. 39—40; T. 28; I. 9—10.

Varietas a planta typica differens lobulis lateralibus semicellulae multum reductis, marginibus lateralibus paululum invaginatis magis quam incis; spinescentia lobulorum polarium necnon reductis; polus a latere visus truncatus, protuberatione utroque in latere, in apice, autem, late convexus. Long. 58—60; lat. 39—40; crass 28; isthm. 9—10.

Florida: Ditch 1 mile west of Holopaw, Osceola Co. (Fla. 135).

Swamp 15 miles west of Sanford, Volusia Co. (Fla. 138).

Pond 3 miles west of Sorrento, Lake Co. (Fla. 140).

Pond 8 miles east of Myakka State Park, Sarasota Co. (Fla. 167).

*Euastrum fissum* West & West var. ANGUSTUM var. nov. Pl. III, Fig. 7.

Cells proportionately narrower than in the typical, about twice as long as wide; cells nearly rectangular in outline, widest in the region just above the base of the semicells, the width reduced at the polar lobe to about that of the basal lobe at the isthmus; with supernumerary granules within the margins of the lobules of the polar lobe; in side view similar to the typical, but with a granule on each side just below the apex. L. 38—41; W. 19—20; T. 15; I. 5.

Cellulae proportione angustiores quam in planta typica, saltem duplo longiores quam latae; cellulae ambitu fere rectangulares; latissimae regione admodum supra basim semicellularum, latitudo ad lobum polarem ad circa latitudinem lobi basalis ad isthmum reducta; granula supranumeraria intra margines lobulorum lobi

apicalis; semicellula a latere visa plantae illi typicae similis, praebens, autem, granulum utrimque admodum infra apicem. Long. 38—41; lat. 19—20; crass. 15; isthm. 5.

Florida: Swamp 9 miles southeast of Naples, Collier Co. (Fla. 80).

Swamp 14 miles west of Torrey, Hardee Co. (Fla. 114).

Ditch 9 miles east of Bradenton, Manatee Co. (Fla. 121).

Swamp 7 miles west of Melbourne, Brevard Co. (Fla. 131)

*Euastrum fissum* West & West var. DECORATUM var. nov.  
Pl. I, Figs. 1, 2.

A variety twice the diameter in length, narrower at the poles than at the base of the semicells, the basal angles rather sharply rounded and furnished with a short tooth-like spine, the lateral margins at first converging and then diverging to form the polar lobe, the upper margin of which slopes toward the relatively deep and narrow apical incision, the tip of the polar lobules sometimes furnished with sharp granules; the upper lateral angles and margins of the polar lobe devoid of spines characteristic of the typical; face of the semicell decorated with a central area of granules arranged in three vertical series of two or three; a protuberance in the mid-region of each apical lobule furnished with a circular arrangement of rather large granules; a granule just within the upper margin of the apical lobules, and two or three granules within the margin of the basal lobes of the semicell; a granule on either side of the apex of the polar notch; sinus narrow and closed throughout; in lateral view semicell triangularly cone-shaped in outline, the basal angles and the apex broadly rounded, with a series of three prominent granules showing along the margin of the base of the semicell, a granule on each margin about half-way to the apex above which is a prominent cylindrical protuberance tipped with three granules, with a marginal granule on each side just above the protuberance and just below the apex which is smooth. L. 54—59; W. 27—30; T. 18; I. 7—9.

Varietas duplo longior quam lata, angustior ad polos quam ad basim semicellularum; angulis basalibus quasi acute rotundatis et spina brevi dentiformi praeditis, marginibus lateralibus convergentibus, deinde, ad lobum polarem efficiendum, divergentibus, margine superiore lobi polaris ad incisionem apicalem, relative profundam angustamque devecto, cacumine lobulorum polarium granulis acutis interdum praedito; margines laterales superiores atque anguli lobi polaris carentes spinis plantae typicae propriis; superficies semicellulae area centrali granulorum, in 3 seriebus verticalibus binorum aut trinorum ordinatorum, ornata; protuberatio media in parte cuiusque lobuli apicalis ordinatione circulari granulorum satis magnorum praedita; granulum unum admodum intramarginem

superiorem lobulorum apicalium atque 2 vel 3 granula intra marginem loborum basaliū semicellulae; granulum unum utroque in latere apicis incisurae polaris; sinus angustus, prorsus inapertus; semicellula a latera visa ambitu triangulariter conica, angulis basalibus atque apice late rotundatis, serie 3 granulorum prominentium secundum marginem basis semicellulae visa, granulo uno utroque in margine, media in parte; apex protuberationem cylindricam prominentem, in cacumine 3 granulis munitam, formans, granulo marginali utrimque admodum supra protuberationem atque infra apicem levem. Long. 54—59; lat. 27—30; crass. 18; isthm. 7—9.

Florida: Ditch 5 miles west of Wewahitchka, Gulf Co. (Fla. 48).

This variety is larger than the typical or any of the described varieties, and is somewhat narrower in proportions. It has much in common with *E. elegans* (Bréb.) Kuetz. and some of its numerous expressions. The pincers type of polar lobe with a deep apical notch identify the Florida specimens with the *E. fissum* series rather than with *E. elegans*.

*Euastrum fissum* West & West var. PSEUDOELEGANS var. nov. Pl. I, Fig. 3.

A variety larger than the typical, about twice the diameter in length; semicells wider at the apex than at the base; basal angles sharply rounded, the lateral margins converging throughout two-thirds of the distance to the apex and then diverging to the upper lateral angles which bear a straight, horizontally projecting spine: the dorsal margin of the apical lobe sloping gently to the deep median notch and forming a broadly rounded apex; basal angles with a short blunt tooth on the margins and with three granules in a triangular pattern just within the margin; with a very slight protuberance of the margin about half-way to the apex; face of semicells with a protuberance in the mid-region furnished with three vertical rows of granules, three in each row; with a protuberance bearing a circular pattern of granules in the mid-region of the two apical lobules and with a bidentate granule just within the margin at the apex on either side of the median notch; sinus linear and closed; lateral view of semicell ovate in outline, lateral margins convex at the base and extending into prominent lateral protuberances bearing three crenations, upper lateral margins deeply retuse between the basal protuberances and a bidentate protuberance on each side just below the apex which is rather sharply rounded, two granules showing on each side between the upper lateral protuberances and the apex; in end view broadly oval with a tricrenate swelling in the mid-region on each side, the poles broadly rounded and showing a pair of subpolar granules at the margin on each side. L. 58; W. 31; W. polar lobe 33; T. 23; I. 10.



Varietas maior quam planta typica, circa duplo longior quam lata; semicellulae latiores in apice quam in basi; anguli basales acute rotundati, marginibus lateralibus per  $2/3$  spatii ad apicem convergentibus, deinde ad angulos laterales superiores, spinem rectam horizontaliter eminentem ferentes, divergentibus; margo dorsalis lobi apicalis ad incisuram mediam profundam leniter devectus et apicem late rotundatum efficiens; anguli basales dente breviter obtuso in marginibus atque 3 granulis in triangulum ordinatis admodum intra marginem praediti; protuberatio minima marginis circa media in parte; superficies semicellularum protuberationem, media in parte 3 ordinibus verticalibus granulorum, 3 utroque in ordine praeditam, habens; protuberatio ordinationem spinarum circularem ferens media in parte lobulorum apicalium duorum, et granulum bidentatum admodum intra marginem in apice utroque in latere incisurae mediae visum; sinus linearis inapertusque; semicellula a latere visa ovata, marginibus lateralibus ad basem convexis et in protuberationes laterales prominentes, 3 crenationes ferentes, extensis, marginibus lateralibus superioribus inter protuberationes basales et protuberationem bidentatum utrimque admodum infra apicem quasi acute rotundatum profunde retusis, granulis duobus utrimque inter protuberationes laterales superiores et apicem visis; semicellula a polo visa late ovata, inflatione tricrenata media in parte utrimque, polis late rotundatis, par granulorum subpolarium in margine utrimque praebens. Long. 58; lat. 31; lat. lobi polares 33; crass. 23; isthm. 10.

Florida: Lily pond 21 miles east of Panama City, Bay Co. (Fla. 43).

This variety is suggestive of *E. elegans* especially because of the broad, almost flat apical margin and the horizontally directed spines at the angles of the polar lobe. The deep polar incision and the form of the cell as seen in lateral view identify it with *E. fissum*.

*Euastrum* FLORIDENSE sp. nov. Pl. I, Fig. 11.

Cells large, a little more than twice the diameter in length, subrectangular in outline with the polar lobe as wide as the basal lobes of the semicell; lower lobules broadly rounded or with margins subparallel, above which is a deep V-shaped invagination of the margin that extends upward to form broadly rounded upper lateral lobules; a deep, narrow sinus separating the lower part of the semicell from the broad and high polar lobe, the upper lateral margins of the polar lobe converging to a truncate apex in which the polar incision is narrow and moderately deep; face of the semicell with one median and two lateral protuberances (one on either side), just above the isthmus and the sinus; wall coarsely scrobiculate; mucilage pore lacking (?); in side view elongate-oval, the poles broadly rounded and smooth, the lateral margins scarcely showing any swellings but

nearly entire, the median incision shallow and slightly open, the sides parallel. L. 104; W. maximum 48; W. at base of semicell 46; T. about 32; I. 16.

Cellulae magnae, paulo magis quam duplo longiores quam latae, ambitu subrectangulares, lobi polari latitudine aequo lobis basalibus semicellulae; lobulis lateralibus inferioribus late rotundatis aut margines subparallelos habentibus; supra, invaginatio marginis profunde, V-formis, ad lobulus laterales superiores late rotundatos formandos, sursum extensa; sinus profundus angustusque partem semicellulae inferiorem a lobo polari lato altoque seiungens, marginibus lateralibus superioribus lobi polaris ad apicem truncatum, incisione polari angusto atque quasi profundo praeditum, convergentibus; superficies semicellulae protuberationem mediam atque duas protuberationes laterales (unam utrimque) admodum supra isthmum sinumque praebens; membrana grosse scrobiculata; porus mucosus deest (?); cellula a latere visa elongato-ovata, polis late rotundatis atque levibus, marginibus lateralibus fere integris, inflationibus quasi nullis, incisione media non alta, paululum aperta, lateribus parallelis. Long. 104; lat. maximum 48; latitudo in basi semicellulae 46; crass. circa 32; isthm. 16.

Florida: Pond 20 miles east of Panama City, Bay Co. (Fla. 47).

This rare species is nearest *E. Allenii* Cushman but differs especially in the form of the polar lobe. *E. Allenii* has the upper lateral margins of the polar lobe with broad and deep invaginations. Also it does not have the narrower and deep sinus between the basal lobes and the polar lobes of the cell. Like *E. Allenii*, however, the Florida plant is broader in the mid-region of the semicell than at the basal lobes.

*Euastrum giganteum* (Wood) Nordst. Pl. I, Fig. 13.

Cells large, elongate, subrectangular in outline, the margins of the cells subparallel, the poles broadly truncate with rather abruptly rounded angles, lateral margins entire, slightly convex from the narrow, deep sinus to the truncate poles, just below which the margins are slightly retuse; median notch narrows relatively shallow; face of semicell with a large supraisthmian protuberance with a lower protuberance on each side just above the isthmus; in side view cells elongate, about the same shape as in front view, but about one-eighth less in thickness than in width, the poles broadly rounded, margins entire and subparallel. L. 202—218; W. 66—70; T. 59; I. 25—29.

Florida: Pond 2 miles west of Milton, Santa Rosa Co. (Fla. 66).

*Euastrum hypochondrum* Nordst. fa. DECORATUM fa. nov. Pl. III, Fig. 10.

A form similar to the typical but larger in all dimensions, and with the median facial protuberance more prominent, the wall throughout decorated with larger granules which are round or conical; in end view usually with a relatively deep invagination of the wall on either side of the median protuberance. L. 60—72; W. 57—66; T. 35—39; I. 12—15.

Forma plantae typicae similis, maior, autem, ubique, necnon protuberatione faciali media magis prominente, tota membrana granulis maioribus, rotundis conicisve ornata; semicellula a vertice visa invaginationem membranae relative profundam utroque in latere protuberationis mediae plerumque praebens. Long. 60—72; lat. 57—66; crass. 35—39; isthm. 12—15.

Florida: Lake near Lebanon Station, Levy Co. (Fla. 74).

Ditch 2 miles south of Bonita Springs, Lee Co. (Fla. 79).

Pond 21 miles southwest of Ocala, Marion Co. (Fla. 163).

*Euastrum hypochondrum* Nordst. fa. PROMINENS fa. nov.  
Pl. III, Fig. 13.

A form in which the central facial protuberance is narrower and more prominent as seen in side or end view than in the typical, the lower lateral margins of the basal lobes somewhat more divergent than in the typical, with three rows of granules encircling the polar lobes. L. 54—56; W. 49—55; T. 25—28; I. 11—12.

Forma protrusione centrale faciali, a latere verticeve visa, angustiore et prominentiore quam in planta typica, marginibus lateralibus inferioribus loborum basaliu aliquanto magis divergentibus quam in planta typica; tres ordines granulorum lobos polares circumdantes. Long. 54—56; lat. 49—45; crass. 25—28; isthm. 11—12.

Florida: Ditch 7 miles south of Fort Myers, Lee Co. (Fla. 78).

Ditch 8 miles south of Punta Gorda, Charlotte Co. (Fla. 77)

Swamp 9 miles south of Naples, Collier Co. (Fla. 80).

Pool in coral rock, 6 miles southwest of Homestead, Dade Co. (Fla. 84).

*E. hypochondrum* and its variations show definite relationships with *E. platycerum* Reinsch, *E. madagascariense* (West & West) Krieg., and a group of species which have no polar notch, all of which are so distinctive that they present a very clear evolutionary series, ending with *E. verrucosum* Ehrbg., or beginning with it, depending on one's point of view in respect to reduction of the cell complexity in the evolution of desmids.

*Euastrum informe* Borge fa. OCULATUM fa. nov. Pl. III, Fig. 8.

A form differing from the typical by having a large mucilage pore medianly located just above the facial protuberance of the semicell. L. 39—44; W. 18—19; T. 14; I. 5.



Forma a planta typica differens possessione pori mucosi magni media in parte admodum supra protuberationem facialem semicellulose. Long. 39—44; lat. 18—19; crass. 14; isthm. 5.

Mississippi: Pond, Highway 26, 3 miles west of Pascagoula River Ferry, George Co. (Miss. 74).

Florida: Lily pond 13 miles east of Fort Walton, Okaloosa Co. (Fla. 6).

Lily pond 21 miles west of Panama City, Bay Co. (Fla. 43)

Pond 3 miles west of Sorrento, Lake Co. (Fla. 140).

This is the third record only of the occurrence of this species, found first in Brazil by Borge, later (1935) in Massachusetts by Prescott. The Massachusetts specimens definitely were without mucilage pores, but whether this is true also of the type specimen cannot be determined inasmuch as Borge was not able to give a full description of the wall features. In the Mississippi and Florida specimens, it is very difficult to determine whether the pore is present in the case of living plants, or preserved cells that retain the chloroplast. It is present in all of the empty cells or semicells that have been examined from these States.

*Euastrum insigne* var. *lobulatum* Presc. & Scott fa. ACUTILOBUM fa. nov. Pl. II, Fig. 4.

A form differing from the typical variety by having the lower basal angles of the semicell narrower and much extended horizontally, and by having prominent lobules on the upper margins of the basal lobes; the polar lobe narrower with subparallel lateral margins, the apical lobules much produced and horizontally extended; sinus more open with widely divergent sides; wall deeply scrobiculate. L. 111; W. 78; I. 15.

Forma a varietate typica differens angulis basalibus inferioribus semicellulae angustioribus et horizontaliter multum extensis, differens necnon possessione lobulorum prominentium in marginibus superioribus loborum basaliu; lobus polaris angustior, marginibus lateralibus subparallelis, lobulis apicalibus multum productis atque horizontaliter extensis; sinus apertior, lateribus late divergentibus; membrana profunde scrobiculata. Long. 111; lat. 78; isthm. 15.

Florida: Pond 6 miles south of Masaryk, Hernando Co. (Fla. 110).

*Euastrum* INVAGINATUM sp. nov. Pl. I, Fig. 8.

Cells medium-sized, about twice the diameter in length; narrowly oval in outline; the semicells semi-oval in shape with a very deep polar notch that is broadly open internally to form a prominent invagination that extends almost to the mid-region of the semicell, the resulting lobules of the cell pincer-shaped; margin of the semicell

showing about nine crenulations between the sharp basal angles and the apex; face of the semicell with an almost imperceptible swelling in the mid-region; the wall thin and coarsely punctate; median incision deep, the sinus narrow and closed throughout; in lateral view the cell fusiform in outline with a V-shaped incision in the mid-region, the apices truncate, the base of the semicell only slightly swollen, the margins undulate; in end view quadrate-oval, the lateral margins nearly parallel, with a slight protrusion on each side, the poles broadly convex and crenate, with three or four series of crenations within the margin. L. 42; W. 20—25; T. about 12; I. 7—8.

Cellulae mediocres, circa duplo longiores quam latae; anguste ovatae; semicellulae semi-ovatae, incisura polari profundissima, ad invagationem prominentem, prope ad mediam semicellulam extensam, formandam, interne late aperta, lobulis cellulae sic formati forcipiformibus; margo semicellulae circa 9 crenulationes inter angulos basales acutos atque apicem praebens; superficie semicellulae inflationem vix perceptam media in parte habens; membrana grosse punctata; incisio media profunde, sinu angusto et prorsus inaperto; cellula a latere visa fusiformis. incisionem V-formem media in parte praebens, apicibus truncatis, basi semicellulae paululum inflata, marginibus undulatis; cellula a polo visa quadrato-ovata, marginibus lateralibus fere parallelis, protrusione minore utrimque media in parte, polis late convexis crenatisque, 3 vel 4 series concentricas crenationum intra marginem praebentibus. Long. 42; lat. 20—25; crass. circa 12; isthm. 7—8.

Florida: Ditch 3 miles east of Gulf Breeze, Santa Rosa Co. (Fla. 2).

Ditch 0.5 mile south of Westbay, Bay Co. (Fla. 9).

This delicate little desmid has a combination of characteristics that make it unlike any known species of *Euastrum*, though there is a distant resemblance to *E. bilobum* Lutkem., which according to Krieger (1937, p. 109) is an arctic species. The marginal crenulations shown in front view combine with those of the vertical view to form a "rippled" surface, which is only noticeable in oblique views. The cell is so thin as to give a superficial resemblance to *Tetmemorus* when seen in side view, and suggests a possible intermediate form.

*Euastrum pectinatum* Bréb. var. **LOBULIFERUM** var. nov.  
Pl. I, Fig. 12.

This variety differs from the typical by having thicker lateral lobules of the polar lobe of the semicells, their apices more broadly rounded, and in having thicker, bilobate lower lateral lobes; the sinus narrow and closed throughout; also differing by having a supracentric mucilage pore in the face of the semicell surrounded by three low swellings, and with an intramarginal mucilage pore on

each side of the polar incision which is characteristically very shallow; wall finely scrobiculate; in lateral view semicell thicker than in the typical, with marginal protuberances just below the poles rather than an invagination, also with secondary lobules at the base of the semicell where the median incision is narrow and deep; in end view broadly elliptic in outline, with two prominent swellings in the mid-region on each side, and with two lower swellings between the mid-region and the poles (six marginal swellings on each side). L. 70—74; W. 52—54; T. 30; I. 11—12.

Varietas a planta typica differens lobulis lateralibus lobi polaris semicellularum crassioribus, apicibus lobulorum latius rotundatis, differens necnon lobis lateralibus inferioribus bilobatis crassioribus; sinus angustus atque prorsus inapertus; differens necnon possessione pori mucosi supracentri in semicellulae superficie, tribus incrassationibus inferioribus circumdati, possessione necnon pori mucosi intramarginalis utroque in latere incisionis polaris, proprie quidem minime alti; membrana subtiliter scrobiculata; semicellula a latere visa crassior quam in planta typica, praebens admodum infra polos protuberantiones marginales potius quam invaginationem, praebens necnon lobulos secundarios in basi semicellulae, ubi incisio media angusta profunda est; semicellula a vertice visa late elliptica, 2 inflationes prominentes utrimque media in parte atque 2 inflationes inferiores inter partem mediam et polum habens (6 inflationes marginales utrimque). Long. 70—74; lat. 52—54; crass. 30; isthm. 11—12.

Florida: Ditch 0.5 mile south of Cottondale, Jackson Co. (Fla. 31).  
Ditch 17 miles north of Okechobee, Okechobee Co.  
(Fla. 91).

*Euastrum pingue* Elfv. Pl. I, Fig. 7.

Cells medium sized, about  $1\frac{1}{2}$  times longer than broad, oval in outline; semicells truncate-pyramidal, the basal angles broadly rounded, the upper lateral margins deeply retuse above the broad basal angles, and then approximately straight to form the polar lobe which is truncate or broadly convex; apical notch narrow and sharp but relatively shallow; face of semicell with a broad, low swelling eccentrically placed to the left of the vertical center-line in the upper semicell and to the right of the center-line in the lower semicell, with a prominent supramedian mucilage pore also eccentrically placed to the right of the center-line in the upper semicell and to the left in the lower one; with a prominent supraisthmian granule at the base of the semicell; wall finely scrobiculate, especially at the angles and on the median protuberance; in side view semicells inversely subnapiform, the basal portion broadly oval to nearly circular in



outline, but then narrowed with retuse margins to form a convex pole, a prominent granule showing at the base of the semicell on each side, and a mucilage pore well within the left margin of the upper semicell about half-way to the pole, and a similar pore within the righthand margin of the lower semicell, the swollen mid-region showing a series of crenulations at the margin; in end view quadrato-oval, the poles truncately convex, the mid-region with a prominent crenulate protuberance which is diagonally opposite the protuberance on the other side, a mucilage pore showing just within the margin midway between the poles and the central protuberance, diagonally opposite a pore on the other side. L. 50—55; W. 36—38; T. 25; I. 12.

Florida: Lily pond 11 miles east of Fort Walton, Okaloosa Co. (Fla. 5).

Previous illustrations of this desmid do not clearly bring out the fact that the central swellings as well as the mucilage pores are eccentrically placed with regard to the vertical center-line. Also it is to be noted that the illustration in Krieger (1937, Pl. 61, Fig. 22) has been reversed, no doubt in the copying process, so that the mucilage pores are shown on the wrong side of the center-line.

*Euastrum platycerum* var. *acutibulum* Krieg. fa. DENTIFERUM fa. nov. Pl. III, Fig. 12.

A form differing from the typical variety by having a more highly decorated wall in the form of granules both on the face of the semicell and around the poles, and by having a prominent tooth-like supraisthmian granule; the lateral lobes of the semicells more nearly horizontal than in the typical variety. L. 57—60; W. 54—56; T. 27; I. 11—12.

Forma a varietate typica differens membrana magis ornata, granulis et in superficie semicellulae et circum polos visis, differens necnon possessione granuli prominentis dentiformis supraisthmalis; lobi laterales semicellularum plus horizontales quam in varietate typica. Long. 57—60; lat. 54—56; crass. 27; isthm. 11—12.

Florida: Swamp 9 miles southeast of Naples, Collier Co. (Fla. 80).

*Euastrum sphyroides* Nordst. fa. GRANULATUM fa. nov. Pl. III, Fig. 11.

A form differing from the typical by having three rows of granules encircling the apical lobe (one row intramarginal), and in having a prominent supraisthmian granule just below the facial protuberance which is decorated with a central pattern of four granules, encircled by a ring of granules in which the one nearest the isthmus is larger. L. 52—54; W. 43—51; T. 22—26; I. 10—12.

Forma a planta typica differens possessione 3 ordinum granulorum

apicalem circumdatum (ordine uno intramarginali), atque granuli supraisthmalis prominentis admodum infra protuberationem facialem, ordinatione centrali 4 granulorum ornatam, hac annulo granulorum, illo, quod isthmo proximum, maiore, circumdata. Long. 52—54; lat. 43—51; crass. 22—26; isthm. 10—12.

Florida: Ditch 18 miles south of Venice, Sarasota Co. (Fla. 75).  
Pond in coral rock, 6 miles southwest of Homestead, Dade Co. (Fla. 84).

*Euastrum Turneri* West & West fa. PORIFERUM fa. nov. Pl. III, Fig. 6.

A form differing from the typical by its possession of two mucilage pores in a horizontal line just above the facial protuberance of the semicell, and by having a much deeper polar incision; the wall lightly but rather irregularly scrobiculate; in lateral view without the undulating upper lateral margins characteristic of the typical. L. 46—51; W. 31—36; T. 19; I. 7—9.

Forma a planta typica differens possessione duorum pororum mucosorum horizontaliter ordinatorum admodum supra semicellulae protuberationem superficialem, differens necnon incisione polari multo profundiore; membrana subtiliter, paululum irregulariter autem, scrobiculata; semicellula a latere visa sine marginibus superioribus undulatis propriis plantae typicae. Long. 46—51; lat. 31—36; crass. 19; isthm. 7—9.

*Euastrum Turneri* var. *stictum* (Boerges.) nobis comb. nov. (Syn. *E. denticulatum* var. *stictum* Boerges.). Pl. I, Fig. 9.

This desmid has been found at three stations in Florida, and also in material from Arnhem Land in North Australia which we have examined (to be published later). Its characteristics agree very well with the original illustration of the Brazilian plant by Boergesen (1890), and also with the figure of the Queensland specimens by Borge (1911). The North Australian collections contain a number of desmid species that were found by Borge in the adjacent State of Queensland, so that it seems quite likely that the Arnhem Land specimens are the same as those seen by Borge. Boergesen's assignment to *E. denticulatum*, however, seems incorrect, since there are too many differences between the plants. Krieger (1937) lists Boergesen's name as a synonym of *E. Turneri* W. West. This also seems undesirable, because of the much deeper apical notch, and the smoothly undulate lateral margins (with projecting granules) as shown in our figure, as compared with the illustrations by West & West (1905) Pl. XXXVII, Figs. 9, 10, which show a deep incision separating the apical lobules from the lateral lobes, and the bilobulate lateral lobes

noted in the description by West & West. We think the best solution of the difficulty is to name the plant as a variety of *E. Turneri*, using Boergesen's varietal name, var. *stictum*. L. 36—37; W. 25—26; T. about 18; I. 7.

Florida: Pond 5 miles south of Marianna, Jackson Co. (Fla. 20).

Swamp 7 miles south of Greenville, Madison Co. (Fla. 238).

Ditch 8 miles east of Hosford, Liberty Co. (Fla. 245).

*Euastrum ventricosum* Lund. var. *SOPCHOPPIENSE* var. nov.  
Pl. II, Fig. 2.

Cells large, 1.8 to 2 times as long as wide; basal angles sharply rounded and extended so as to overlap the angles of the other semicell, thus closing the median sinus which is deep and enlarged internally; lateral margins broadly concave and then diverging into the upper lateral lobules which are narrow and directed upward at a sharp angle, with a deep sinus between them and the polar lobe which is broadly cuneate; the apical margin convex and with a narrow polar incision; face of the semicell with a prominent supraisthmian protuberance and with a smaller intramarginal protuberance on either side at the base of the semicell, and with a low protuberance in the mid-region of the apical lobules; wall coarsely scrobiculate; in lateral view semicell truncate-pyramidal with the margins evenly converging to an enlarged apex which is trilobed; vertical view ellipsoid in outline with a protuberance on either side of the sharply rounded poles and a prominent lateral protrusion in the mid-region. L. 169—180; W. 91—99; T. 61—64; I. 23—25.

Cellulae magnae, 1.8—2.0 plo longiores quam latae; anguli basales acute rotundati et ad angulos semicellulae alterius obtendendos extensi, hoc modo sinum medium, profundum et interne ampliatum, claudentes; margines laterales late concavi, deinde in lobulos laterales superiores divergentes, his angustis et abrupte sursus directis; sinu profundo inter hos et lobum polarem late cuneatum; margo apicalis convexus, incisione polari angusta; superficies semicellulae protuberatione media surpaisthmali prominente, atque protuberatione intramarginali minore utrimque in basi semicellulae atque protuberatione humili media in parte lobulorum apicalium praedita; membrana grosse scrobiculate; semicellula a latere visa truncato-pyramidalis, marginibus ad apicem amplificatum, trilobatum aequaliter convergentibus; semicellula a vertice visa ellipsoidea, protuberationem utroque in latere polorum acute rotundatorum, atque protrusionem lateralem prominentem media in parte praebens. Long. 169—190; lat. 91—99; crass. 61—64; isthm. 23—25.

Florida: Pond 3 miles south of Sopchoppy, Wakulla Co. (Fla. 154).



This variety differs from the typical mostly in the deeper and inwardly opening sinus between the polar lobe and the base of the semicell, and in the narrowness and sharpness of the basal angles. Also the Florida plants have no central mucilage pore nor the protuberances on either side of it which characterize the typical. It approaches some of the varieties of *E. crassum* (Bréb.) Kuetz., with which it should be compared.

*Euastrum verrucosum* Ehrbg. fa. Pl. II, Fig. 5.

In this form the polar lobe is more produced than in the typical because the upper lateral sinuses are deeper and narrower and hence the upper lateral lobules are relatively narrower and longer. L. 82; W. 68; I. 19.

Florida: Pond 22 miles southwest of Ocala, Marion Co. (Fla. 164).

*Euastrum verrucosum* var. *alatum* Wolle fa. EXTENSUM fa. nov. Pl. II, Fig. 6.

A form differing from the typical by having narrower basal lobules, which are longer and more nearly horizontally directed; the apical lobe broad, with the upper margin less retuse; granulations similar to the typical except that there are two prominent granules at the apex and a definite row of granules across the base of the semicell just above the isthmus. L. 85—87; W. 79—88; T. 42; I. 18.

Forma a varietate typica differens lobulis basalibus angustioribus, longioribus et potius horizontaliter extensis; lobus apicalis latus, margine superiore minus retuso; granulationes plantae typicae similes nisi quod habet duo granula intramarginalia prominentia in apice atque ordinem definitum granulorum trans basim semicellulose admodum supra isthmum. Long. 85—87; lat. 79—88; crass. 42; isthm. 18.

Florida: Swamp 9 miles southeast of Naples, Collier Co. (Fla. 80).

#### REFERENCES

- BOERGESEN, F. - 1890. - Desmidiaceae in: E. Warming, Symbolae ad floram Brasiliae centralis cognoscendam. Vid. Meddel. fra d. naturh. Foren. 1890; 929—958.
- BORGE, O. - 1911. - Algologische Notizen 6. Süßwasseralgen aus Queensland. Bot. Notis. 1911; 197—206.
- KRIEGER, W. - 1937. - Die Desmidiaceen, in Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz. Band 13, Teil 1.
- PRESCOTT, G. W. and SCOTT, A. M. - 1945. - The Fresh-Water Algae of Southern United States III. The Desmid Genus *Euastrum*. Am. Midl. Nat. 34 (1): 231—257.

## EXPLANATION OF PLATES

### PLATE I.

- Figs. 1, 2. *Euastrum fissum* West & West var. *decoratum* var. nov.  
3. *E. fissum* West & West var. *pseudoelegans* var. nov.  
4. *E. bidentatum* Naeg. var. *quadrioculatum* var. nov.  
5. *E. Ciastonii* Racib. var. *apertisinuatum* var. nov.  
6. *E. Ciastonii* Racib. var. *asymmetricum* var. nov.  
7. *E. pingue* Elfv.  
8. *E. invaginatum* sp. nov.  
9. *E. Turneri* var. *stictum* (Boerges.) nobis, comb. nov.  
10. *E. acmon* West & West var. *turgidum* var. nov.  
11. *E. floridense* sp. nov.  
12. *E. pectinatum* Bréb. var. *lobuliferum* var. nov.  
13. *E. giganteum* (Wood) Nordst.

### PLATE II.

- Fig. 1. *Euastrum crassum* (Bréb.) Kuetz.  
2. *E. ventricosum* Lund. var. *sopchoppiense* var. nov.  
3. *E. crassum* var. *scrobiculatum* Lund. fa.  
4. *E. insigne* var. *lobulatum* Presc. & Scott fa. *acutilobum* fa. nov.  
5. *E. verrucosum* Ehrbg. fa.  
6. *E. verrucosum* var. *alatum* Wolle fa. *extensum* fa. nov.

### PLATE III.

- Fig. 1. *Euastrum bidentatum* var. *oculatum* (Istv.) Krieg. fa. *glabrum* fa. nov.  
2. *E. evolutum* var. *monticulosum* (Tayl.) Krieg. fa. *poriferum* fa. nov.  
3. *E. evolutum* var. *reductum* var. nov.  
4, 5. *E. evolutum* var. *integrius* West & West fa. *turgidum* fa. nov.  
6. *E. Turneri* W. West fa. *poriferum* fa. nov.  
7. *E. fissum* West & West var. *angustum* var. nov.  
8. *E. informe* Borge fa. *oculatum* fa. nov.  
9. *E. denticulatum* var. *rectangulare* West & West fa.  
10. *E. hypochondrum* Nordst. fa. *decoratum* fa. nov.  
11. *E. sphyroides* Nordst. fa. *granulatum* fa. nov.  
12. *E. platycerum* var. *acutilobum* Krieg. fa. *dentiferum* fa. nov.  
13. *E. hypochondrum* Nordst. fa. *prominens* fa. nov.

Addresses of the authors:


A. M. SCOTT

2824 Dante Str., New Orleans La.

G. W. PRESCOTT


Department of Botany and Plant pathology  
State College, East Lansing Michigan

# I






## II



# III



# On the nuclear cytology of *Hydrodictyon*

by

JOHANNES PROSKAUER

In the autumn of 1948 Dr. M. A. POCKOCK suggested to me a cytological investigation of three members of the genus *Hydrodictyon*, on which she has been carrying out extensive studies.

In the literature on the genus there are a number of descriptions of nuclear phenomena. TIMBERLAKE (1901) gave a detailed account of nuclei and their division in *H. reticulatum*. Because of the small size of the chromosomes he was unable to achieve an accurate count, but mentioned that, in a few instances, he could discern ten. He also noted certain apparent anomalies. YAMANOUCI (1913) furnished a description of mitosis in his *H. africanum*, where he counted eighteen chromosomes. My observations on mitosis confirm and broaden the general conclusions of these workers, but no centrosomes or spindle fibres were seen, and the splitting of the spindle at anaphase figured by YAMANOUCI appears to lack basis in fact. As TIMBERLAKE had already concluded, mitosis in these organisms is essentially 'normal'. TIMBERLAKE's and YAMANOUCI's papers show that good information on nuclear phenomena in algae is not just a development of recent years.

MAINX (1931) noted in passing that he had observed typical meiotic synapses in germinating zygospores of *H. reticulatum*. He concluded that the species was a typical haplont, as shown below on somewhat more ample evidence.

The paper by PALIK (1949) on *H. reticulatum*, therefore, came as a considerable surprise. So-called division of large stellate 'nuclei' was described at length, and he concluded that the process was markedly different from normal mitosis, and unlike anything before, seen in plants. In addition meiosis was stated to take place during gametogenesis, the haploid chromosome complement being two. Examination of his many photographs makes it evident that the large stellate bodies, the division of which was described, were not nuclei. Where the former are shown, the latter do not appear to have stained. The most probable explanation is that the stellate bodies represent



localized fixative or other mineral precipitates. Unfortunately it is not indicated which one of several described techniques was employed in the preparation of particular specimens.

The description of meiosis, however, was based on observations on real nuclei as shown by the illustrations, but these nuclei appear to have been undergoing normal mitosis. The chromosome counts were apparently obtained from early anaphases seen in side view, a whole chromosome plate seen edgewise being interpreted as one or two chromosomes respectively. This seems to explain the enucleate cells with postulated nuclear migration into them, and numerous other aberrant phenomena described in the paper.

The following observations are put on record to reemphasize the essentially normal character of the nuclear behaviour. All the described species of *Hydrodictyon*, with the exception of the *H. indicum* of IYENGAR, which is of uncertain standing (cf. POCK, 1937), were available for study and proved uniform in matters of detail.

## MATERIALS

The material, with one exception, was collected by Dr. POCK, whose efforts made this paper possible. My thanks are due to her also for reading the manuscript. The species examined were:

*Hydrodictyon reticulatum* (L.) LAGERHEIM (syn. *H. utriculatum* ROTH).

(1) Recreation Grounds, Fremont, Nebraska, U.S.A. In rainwater pools.

(2) Cherry Creek, Nevada County, California, U.S.A. Collected by Dr. MALCOLM NOBS.

Cultures of nets and zygospores were kindly supplied by Dr. POCK. The nets were cultured in a solution prepared according to the recipe given by JULLER (1937), with added soil extract. Zygospore cultures were obtained by plating out gametes in drops on the surface of agar plates containing the same minerals ( $3/4$  % agar made up in the same culture solution).

*Hydrodictyon patenaeforme* POCK.

(1) Grahamstown, Cape Province, South Africa.

(2) De Klip, Cape Flats, near Cape Town, South Africa. (Type area).

Cultured from soil samples by Dr. POCK.

*Hydrodictyon africanum* YAMANOUCHI.

*Isoetes vlei*, Cape Flats, near Cape Town, South Africa. (Type area)

The plants appeared in a culture from a soil sample supplied by Dr. POCKOCK. This series of cultures was initially made for the purpose of growing *Riella purpureospora*; in addition *Nitella praeclara* and *Isoetes* sp. were present.

## METHODS

The material was fixed in one part ethanoic acid to three parts of ethanol throughout.

(a) Mitosis — Whole mounts. Stained either (a) with Harris' haematoxylin with cytoplasmic counterstain of orange G, phloxine or fast green, or (b) with acetocarmine (see below), with or without cytoplasmic counterstain.

(b) Mitosis — Squashes. Late in the afternoon or early in the morning some cells of a healthy growing net generally showed division. Parts of such nets were fixed and stained by the acetocarmine method described by GODWARD (1948). This involves short washing in water, mordanting up to a minute in 4 % aqueous ferric ammonium alum, short washing in water, followed by staining for a few minutes in saturated acetocarmine. It was found efficacious to open up some of the larger coenocytes after fixation, the material being then mounted in acetocarmine and slightly warmed. This cleared the cytoplasm and tended to decolourize the pyrenoids and to bring out the nuclei and division stages. Any coenocyte showing division was now easily recognised. If large enough, it was cut open with fine scissors, and the protoplast removed with fine needles. It also proved satisfactory merely to cut up the coenocyte into small pieces consisting of wall material with adhering protoplasm. The pieces were remounted in acetocarmine, or, if some destaining was desirable, in 45 % ethanoic acid. The heated preparation was placed on a table and the coverslip pressed down sharply with a cloth-covered finger. The protoplasm tended to spread on the coverslip. The preparations were studied at this stage. They could be made permanent by inverting the slides in dilute ethanoic acid, when the coverslip, with the adhering protoplasm, drops off; after dehydrating the preparation is mounted in 'Diaphane' ('Euparal').

(c) Meiosis. Agar cultures of green zygospores in active growth were employed, i.e. cultures in which the zygospores had not entered the resting phase characterized by production of orange pigment. The agar with the spores on its surface was cut into small blocks which were transferred to 'Pyrex' glass dishes containing 50 % culture

solution or merely glass-distilled water. After one or two days the first zoospores and polyhedra appeared in the solution. When germination was in progress, the blocks were fixed and stained *in toto* by the acetocarmine method described above. The excess agar was trimmed off, and the rest of the block mounted on a slide in acetocarmine. Warming dissolved the agar, leaving the zygospores. Those which had not started to germinate tended to retain some pigment and stained tardily, but repeated heating proved helpful. Changes in the spore membranes during germination allowed better diffusion. The nuclear details were seen more clearly when the preparation was squashed. This liberated the contents from the spore wall and spread them out (fig. 23).

## OBSERVATIONS

Young cells of nets of all the species of *Hydrodictyon* under review start with the single nucleus of the parent spore, and possess one pyrenoid. Nuclei and pyrenoids increase in number as the cells enlarge. I do not desire to enter into the controversy about the chloroplast, except to state that I have been unable to convince myself of its existence as a distinct entity in the older coenocytes. The structure of the mature protoplast appears indistinguishable from that of *Protosiphon*, if a comparable portion of the vesicle of that organism is considered. It also shows some resemblance, albeit somewhat remote, to that of certain species of *Cladophora*.

Nuclear division proceeds independently in different cells of a net; within individual cells there is, however, some synchronization. In smaller coenocytes all the nuclei will be in the same phase of mitosis at the same time; in Fig. 22 all are at metaphase of division. In larger coenocytes nuclear division spreads like a wave. Thus, nuclei at the (variable) point at which division started may be at telophase. These will be surrounded by a belt of anaphases, these in turn by metaphases, prophase, and interphases respectively. Fig. 21 illustrates the range from metaphase to late anaphase in one of the smallest coenocytes in which such division was observed. Ultimately, in large coenocytes, the number of nuclei can generally be estimated to be of the order of tens of thousands, while in *H. patenaeforme* Pocock (1937) estimated it at hundreds of thousands prior to gametogenesis. In such cells some thousand or so nuclei can be found in the same mitotic phase. *Hydrodictyon* may well be suitable for studies on certain aspects of the physiology of mitosis.


The following description of mitosis is based on unsquashed whole mount preparations of *H. reticulatum*, but the details have

been found to be identical in the other two species. The interphase (figs. 1, 10) contains a single central nucleolus; the nucleoplasm appears slightly granular and no definite heterochromatin is observable. In early prophase chromocentres appear in the nucleoplasm (fig. 2). As the chromosomes become recognisable the nucleolus disappears (figs. 3—5). In metaphase the chromosomes form a dense plate. At this stage the spindle is somewhat difficult to demonstrate (figs. 6, 21). Since its axis normally lies parallel to the cell wall, the chromosome plate is seen edge-on in whole mounts and, as the chromosomes are very small and compacted, the plate appears under medium magnification as a thick short line (figs. 21, 22), although under higher magnification individual chromosomes can be distinguished (figs. 24, 28 from squashes). At anaphase the spindle is




Figs. 1—12, 17—20, *Hydrodictyon reticulatum*; 13—14, *H. patenaeforme*; 15—16, *H. africanum* Figs. 1—10, Stages of mitosis from whole mount preparations. Figs. 11—16, Metaphase plates, squashes. Figs. 17—20, Meiosis, squashes of germinating zygospores, fig. 17, probably pachytene, the rest metaphase.  $\times 2400$ .






21




22



23

quite clear (figs. 7, 8, 21), especially between the separating chromosomes. It is, presumably, gelatinous in nature. The ultimate distance of separation is variable. Reorganization at telophase (fig. 9) is difficult to decipher. The sister nuclei (fig. 10) continue to lie close together for some time.

The largest chromosomes do not exceed one micron. In squash preparations the plates flatten out somewhat, but spreading of chromosomes is very limited. The appearance of the plates and chromosomes is identical in all the material of the three species studied, and their number also appears the same. It was impossible to count them with absolute certainty, even with the numerous stages provided by large coenocytes. The number appears to be  $19 \pm 1$ , but most likely 19 or 18. The uncertainty is due to the small size and the possibility of the presence of satellites. Metaphase plate squashes are illustrated for *H. reticulatum* in figs. 11, 12, 25, 26,



Figs. 24—26, 30, 31, *H. reticulatum*; 27—28, *H. patenaeforme*; 29, *H. africanum*. Figs. 24, 28, Mitotic metaphase plates in side view. Figs. 25—29, Mitotic metaphase plates. Figs. 30—31, Meiotic metaphases. All from squashes  $\times 2000$ . (Zeiss Apo. 60 I, 40; Homal iv).

Figs. 21—23. *H. reticulatum*. Fig. 21, Divisions in a small coenocyte; metaphases above, anaphases below.  $\times 1000$ . Fig. 22, Small coenocyte, all the nuclei are at metaphase (the plates appearing as short lines); the round objects are pyrenoids.  $\times 345$ . Fig. 23, Squash preparation of germinating zygospore at meiotic metaphase; note the broken zygospore membrane.  $\times 2000$ .

for *H. patenaeforme* in figs. 13, 14, 27, 28, and for *H. africanum* in figs. 15, 16, 29.

Meiosis could take place in the life cycle of *Hydrodictyon* during the germination of the zygospore or at gametogenesis. The latter possibility is attractive in so far as reduction there might account for the difference in behaviour of zoospores and gametes. No evidence was found to support it. On the other hand, stages of the first nuclear division in the germinating zygospore, the 'orthodox' place, have the aspect of meiotic figures recognised in other plants. Only one nucleolus is generally present in the zygospore nucleus. Detail of prophase stages is, as is common, hard to discern clearly. Fig. 17 shows what is believed to be pachytene. At diakinesis and metaphase the gemini appear considerably larger than the ordinary mitotic chromosomes. In fresh preparations their component chromosomes could sometimes be distinguished. No more than 18 or possibly 19 gemini were observed in any instance, but this does not preclude the possibility that a twentieth may have been hidden somewhere (figs. 18—20, 30, 31). More than two dozen meiotic metaphases were studied.

In the second division within the zygospore, where observed, the chromosomes are of the same size as the ordinary somatic ones. Observations on the fixed preparations suggest that often only two zoospores are formed. There was some vague indication of breakdown of supernumerary nuclei. It remains an open question whether the gametes can also develop parthenogenetically. More work on late stages of germination on living material is needed. Micromanipulation may probably prove to be the most suitable method for investigating the possibility of parthenogenesis.

The observations on meiosis were made on material of *H. reticulatum* only, but it is presumed that the other species show an identical behaviour.

In view of the statements of previous workers, the following abnormality may be mentioned. In the beginning of these studies an attempt was made to induce mitosis by putting cultures of nets into dark electric ovens at slightly elevated temperatures, although this method was not used in any of the work described above. One net of *H. patenaeforme* was placed in an oven at the set temperature of 30° C, but, owing to bad ventilation, the temperature may have risen to somewhat above that figure. In certain cells of this net nuclear divisions with tripolar spindles were observed; those cells which were not fixed were moribund, and died shortly after the exposure. The same phenomenon might occur in nature, where similar temperatures may be reached when a pool dries up. It may explain TIMBERLAKE's tripolar spindles and large irregular nuclei,

as opposed to the large 'nuclei' of PALIK, and may normally occur during the frequently observed death of large populations without reproduction. In squash preparations of normal material a few nuclei with apparently double the normal chromosome complement have been seen. These cases have been interpreted as resulting from squashes of early anaphases, rather than metaphases of diploid nuclei.

## SUMMARY

1) Mitosis was studied in *Hydrodictyon reticulatum*, *H. patenaeforme*, and *H. africanum*; whole mounts and squash preparations being employed. The process is similar to that found in other plants. Division is synchronized, in larger coenocytes it proceeds in wave-form. The behaviour of the different species is identical.

2) Meiosis was observed in germinating zygospores of *H. reticulatum*.

3) The chromosome number is identical in all the species. The number was determined within the limits  $19 \pm 1$ .

4) Elevation of temperature above  $30^{\circ}$  C induced abnormal tripolar nuclear divisions, followed by death.

## ZUSAMMENFASSUNG

1) Die Mitose bei *Hydrodictyon reticulatum*, *H. patenaeforme* und *H. africanum* wurde untersucht, unter Benutzung von Ganz- und Quetschpräparaten. Die Vorgänge weichen von denen in anderen Pflanzen nicht ab. Kernteilung erfolgt synchron (wellenförmig in grösseren Zellen). Die untersuchten Arten verhalten sich vollkommen gleich.

2) Die Reifeteilung wurde bei der Zygotenkeimung von *H. reticulatum* festgestellt.

3) Alle Arten haben die selbe Anzahl von Chromosomen, sie fällt in die Grenzen  $19 \pm 1$ .

4) Erhöhung der Temperatur über  $30^{\circ}$  brachte abnorme tripolare Kernteilungen und darauffolgendes Absterben herbei.



## REFERENCES

- GODWARD, M. B. E. - 1948. - The iron alum acetocarmine method for algae. *Nature* clxi: 202.
- JULLER, E. - 1937. - Der Generations- und Phasenwechsel bei *Stigeoclonium subspinosum*. *Archiv f. Protistenkunde* lxxxix: 55—93.
- MAINX, F. - 1931. - Gametencopulation und Zygotenkeimung bei *Hydrodictyon reticulatum*. *Archiv f. Protistenkunde* lxxv: 502—516.
- MALIK, P. - 1949. - Kernteilung bei *Hydrodictyon urticulatum* Roth. *Index Hort. Bot. Univ. Budapest*. vii: 150—172.
- POCOCK, M. A. - 1937. - *Hydrodictyon* in South Africa. With notes on the known species of *Hydrodictyon*. *Trans. Roy. Soc. South Africa* xxiv: 263—280.
- TIMBERLAKE, H. G. - 1901. - Development and structure of the swarmspores of *Hydrodictyon*. *Trans. Wisconsin Ac. Sc.* xiii: 486—522.
- YAMANOUCHI, S. - 1913. - *Hydrodictyon africanum*, a new species. *Bot. Gaz.* lv: 74—79.

Address of the author:  
DEPARTMENT OF BOTANY  
Berkeley 4, California

## Bibliography

- STUNDL, K. - Zur Hydrographie und Biologie der österreichischen Donau, Schweiz. Ztschr. f. Hydrol., XIII/1, 1951, 36—53, 2 fig., 9 tables.  
On Hydrography and Biology of the austrian Danube. Condensed English summary: The Austrian part of Danube, including some of its tributaries, have been examined for a number of years. Dissolved salts vary very much due to extensive changes in the quantity of water. Pollutions by sewage cause changes in the quantity of ions and reduction in the quantity of oxygen. Some kinds of sewage cause big damage to water organisms and have a bad effect on the stock of fish in some tributaries. It has been found that stream-plankton is swept along from the by-waters to a large extent. Diatoms form the main body of phytoplankton, and rotifers are more frequent than crustacea. In the affluents below Vienna the not purified sewage from the city causes a great increase of organisms in the water and in the mud, which is of great importance for the output of fishery.
- THIENEMANN, A. & K. STRENZKE - Larventyp und Imaginalart bei Chironomus s.s., Ent. Tidsk., 72, 1—2, 1951, 21 p., 12 fig.  
Larval-type and species in the genus Chironomus s.s.; ten species are studied and a key is given.
- ALEEM, A. A. & F. HUSTEDT - Einige neue Diatomeen von der Südküste Englands, Botan. Notiser, 1951, 1, 13—20, 5 fig.  
New Diatoms from the southern coast of England. The following forms are described: Caloneis Hustedti ALEEM, Navicula supralitoralalis AL. & HUST., Amphora tenerrima AL. & HUST., Nitzschia calcicola AL. & HUST., N. valdestriata AL. & HUST., N. salinicola AL. & HUST.
- ALEEM, A. A. & F. HUSTEDT, Littoral Diatoms from the Salstone, near Plymouth, J. Mar. Biol. Ass. U. K., XXX, 1951, 177—196, 2 fig.  
132 species and 9 varieties are mentioned, including the following new species: Cocconeis guttata, Navicula aleemi HUST., N. britannica, Pleurosigma subhyalinum.
- VISSERIJ-NIEUWS, IV, no 3, Juli 1951. - From the contents: Fishery-results. - Rescue-organization on the Dutch coast. - Marked Carp in the IJssel-lake.
- STELLA, E. - Sulla presenza di Cyclops minutus Claus nell'oasi di Gadames, Biologia coloniale, VIII, 1947: 15—21, 3 fig.  
On the presence of Cyclops minutus Claus in the Gadames oases.
- STELLA, E. - Studio biologico preliminare sulle zoocenosi del Tevere, Atti Soc. Intern. Limnol. theor. & appl., XI, 1951: 383—391, with an English summary.  
Preliminary biological study on the zoocenoses of the river Tiber.

- WERFF, A. VAN DER - On a new material for screening plankton, *The Amsterdam Naturalist*, 1, 2, 1950, 53—54, 2 fig.  
Report of tests made with the new Nylon gauze for plankton nets. Although more expensive, it presents many advantages and its use may be recommended.
- WERFF, A. VAN DER - De ontwikkeling van groene algen en andere organismen in plastieke waterleidingbuizen, *Water*, No 1, 4 Jan. 1951, 3 bblz.  
The development of green algae and other organisms in plastics canalization tubes.
- LUTHER, H. - Vorschlag zu einer ökologischen Grundteilung der Hydrophyten, *Acta Bot. Fennica* 44, Helsingfors 1949, 15 p.  
Proposal of an ecological division of the hydrophyta. The division proposed includes four major groups: Haptophyta (Ephaptophyta and Endohaptophyta), Rhizophyta, Planophyta (Planktophyta and Pleustophyta) and Helophyta.
- SYMOENS, J. J. & A. VAN DER WERFF - Note sur les formations du tuf calcaire des environs de Consdorf (Grand Duché de Luxembourg), *Bull. Soc. R. Bot. Belgique*, 83, 1951: 213—218, 1 fig.  
Note on the calcareous tuff formations near Consdorf (Gt. Duchy of Luxembourg). A new species is described: *Lyngbya Vanden Berghemii* Symoens.
- STELLA, E. - Il Lago di Albano, *Boll. di Pesca, Piscicoltura e Idrobiologia*, XXVII, VI (n. s.), 1, 1951, 32 p., 3 fig., 3 graph., 3 tabl., 1 pl.  
The lake Albano, 2d part comprising the littoral communities.
- VISSERIJ-NIEUWS, IV, no 6, Oct. 1951. - Contents: Fishery results. - Annual meeting of the International Council. - Our most important fish species, etc.
- VOLLENWEIDER, R. A. - Eine Schnellmethode zur Ammoniakbestimmung im Tiefenwasser eutropher Seen, *Schweiz. Ztschr. f. Hydrol.*, XIII, 1951, F. 1, 27—28, 1 table.  
A quick method for determination of ammonium in deep water of eutrophic lakes.
- OYE, P. VAN - Au sujet des Rhizopodes du Grand Duché de Luxembourg, *Biol. Jaarb.*, 18, 1951, 82—121, 42 fig.  
On the Rhizopods of the Great Duchy of Luxembourg. The first study ever made on the Rhizopods of that country; 48 forms were collected by the author during a trip of 10 days in May, 1950. Annotated descriptions of all those forms are given; the following are described as new: *Cyclopyxis arcelloides* PENARD, var. *minima* and *Jungia sundanensis* VAN OYE, var. *ovoidalis*. The author points out that the ecological distribution of the Rhizopoda of Luxemburg does not differ from that of the Belgian and Dutch forms, although the relative frequency is nearer to that observed more eastwards. No glaciary form was found.
- IRÉNÉE-MARIE, FRÈRE - Desmidiées de la région de Québec, *Nat. Can.*, 78, 1951, 88—127, 2 pl.  
Desmids of the region of Québec. Description of species belonging

to the genera *Cosmarium* and *Xanthidium*. The following forms are new: *Cosmarium dentatum* WOLLE, var. *glabrum* and *C. vexatum* WEST, var. *quebecense*.

ROHDE, W. - Minor constituents in lake waters, *Proc. Int. Ass. theor. & appl. Limnology*, XI, 1951, 317—323, 7 fig.

VOLLENWEIDER, R. A. - Aspekte moderner Limnologie, Festschr. 131. Jhrvers. Schweiz. Nat.f. Ges. Luzern, 1951, 18 p.  
Aspects of modern limnology. The following points are discussed. Water as "watery solution"; matter exchange; biology of the production.

SAMPAIO, J. - Subsídios para o estudo das Cianofitas das águas termais portuguesas, *Bol. Ass. Filos. Natural*, II, 1951, no 20, 155—160.  
Materials for study of Cyanophyceae of the thermal water of Portugal. Annotated list of 14 species belonging to four genera.

ROSA, K. - Algenflora von Südböhmen, *Studia Botanica Cechoslovaca*, 12, 1951, Fasc. 2, 173—232, 14 pl.  
Algal flora of South Bohemia: I. Algae of the vicinity of Blatná. A list of not less than 398 forms, with description of 9 new ones.

HOOGENRAAD, H. R. - Die fossilen Rhizopoden der Torfstreu, *Biol. Zentralbl.*, 70, 1951, 3/4, 113—119, 2 tables.  
Fossil turf Rhizopoda. Short account on Rhizopods found in subfossil turf; characteristic is the association of 6 thecamoebian Rhizopods viz. *Amphitrema flavum* and *wrightianum*, *Assulina muscorum* and *seminulum*, *Arcella artocrea* and *Bullinula indica*. Some 9 other species are present only sporadically.

HOOGENRAAD, H. R. & A. A. DE GROOT - Thecamoebie Moosrhizopoden aus Südamerika, *Arch. f. Hydrobiol.*, XLV, 1951, 346—366, 20 fig.  
Thecamoebian moss-Rhizopods from South America. Account on a number of Rhizopods found on eight species of moss.

BENTHEM JUTTING, W. S. S. VAN - Dr. Johannes Govertus de Man, Middelburg 2 Mei 1950, Middelburg 19 Januari 1930. Een Zeeuwse zoöloog van internationale vermaardheid, *Biol. Jaarb.*, 18, 1951, 130—250, portr.  
A detailed biographical sketch of the well known crustacea specialist J. G. de Man, of Middelburg.

MARGALEF, R. - Rôle des Entomostracés dans la régénération des phosphates, *Trav. Ass. Int. Limnol. théor. et appl.*, XI, 1951, 246—247, 1 fig.  
The part played by the Entomostraca in the regeneration of the phosphates.

MARGALEF, R. - Observaciones sobre *Triops* (= *Apus*) *cancriformis* de una localidad catalana, *P. Ins. Biol. Apli.*, IX, 1951, 247—254.  
Observations on *Triops* (= *Apus*) *cancriformis* from a locality in Catalonia.



- MARGALEF, R. - Diversidad de especies en las comunidades naturales, *P. Inst. Biol. Apl.*, IX, 1951, 5—27.  
Specific diversity in natural communities.
- BOURRELLY, P. - Une nouvelle espèce de Chrysomonadinae: *Mallomonas Doignonii*, *Bull. de la Soc. bot. Fr.*, 98, 1951, 156—158, 4 fig.  
A new species of Chrysomonadinae: *Mallomonas Doignonii*.
- BOURRELLY, P. - *Euglena Chadefaudii*, une nouvelle espèce d'Euglène viridoïde, *Bull. de la Soc. bot. Fr.*, 98, 1951, 143—145, 1 fig.  
*Euglena Chadefaudii*, a new species of greenish euglenae.
- BOURRELLY, P. et E. MANGUIN - Florule algologique d'une rizière de Camargue, *La Terre et la Vie*, 1950, No 5 (numéro spécial), 286—296, 2 pl.  
Algal florule of a rice-field in the Camargue. List of 144 forms with some remarks.
- BOURRELLY, P. - Protistologica CXII, Note sur les Flagellés incolores, *Arch. Zool. expér. et gén.*, 88, 1951, 73—84, 27 fig.  
Protistologica CXII, Note on the incolorous Flagellata. Genus *Bicoeca* J. CLARK. Description of 3 new species: *B. De Poucquesiana*, *B. Fottii*, *B. stellata*. The author proposes to divide the genus *Bicoeca* in four sections or subgenera, viz. *Eubicoeca*, *Codomonas*, *Stephanocodon* and *Poteriodendron*.
- THIENEMANN, A. - Vom Gebrauch und vom Missbrauch der Gewässer in einem Kulturlande, *Arch. f. Hydrob.*, XLV, 1951, 557—583.  
Of use and misuse of water in a civilised country. The writer emphasizes the wrong sides of using natural waters in economy.
- LHOTSKY, O. - The cell-wall structure in *Closterium moniliferum* (BORY) EHRB., *Vestn. Kral. Ceske Spoll. Nauk*, XII, 1951, 26 p., 8 fig., 2 pl.
- DAMAS, H. - La ponte en aquarium des Lamproies fluviaux et de Planer, *Ann. Soc. Roy. Zool. Belgique*, 81, 1950, 151—162.  
River and Planer's Lampreys laying eggs in aquarium.
- HASLER, A. D. & WISBY, W. J. - Discrimination of stream odors by fishes and its relation to parent stream behavior, *The Amer. Nat.*, 85, no 823, 1951, 223—238, 1 fig., 6 tables.  
Preliminary tests of the theory explaining the mechanism by which migrating salmon returns to their parent streams guided by the characteristic odor of the latter.
- WOHLSCHLAG, D. E. & HASLER, A. D. - Some quantitative aspects of algal growth in Lake Mendota, *Ecology*, 32, no 4, 1951, 581—593, 1 fig., 5 tables.  
With the case history of an individual *Diplocystis* (= *Microcystis*) *aeruginosa* "Bloom".
- SRAMEK—HUSEK, R. - Ciliata-Peritricha Nasich Rek (Prispeoky k poznani ceskoslovenskych nalevniku III), *Casopis Nar. Musea*, CXVII, 1948, 17 pp., 7 pl.

Ciliata-Peritricha of our rivers. Described as new: *Opercularia moldavica* sp. n.

SRAMEK—HUSEK, R. - K poznání planktonu a epizoonu středoevropských rybníků, *Casopis Nar. Musea*, CXVII, 1948, 15 pp., 4 pl.

To the knowledge of plankton and epizoon of Czech ponds. Described as new: *Epicystis chironomi*, *Carchesium ephemerae*, *Vorticella epizoica* and *Vorticella bosminae*.

SRAMEK—HUSEK, R. - O několika zajímavých nálevnicích z Českomoravské vysokiny, *Vestn. Čsl. zool. spol.*, 13, 1949, 325—333, 4 fig.

Some interesting Ciliates of the Czech-Moravian Highlands. Described as new: *Pseudoblepharisma tenue-chlorelligera* var. n. and *Cyclidium sphagnetorum* sp. n.

SRAMEK—HUSEK, R. - Karel Schäferna †, *Proc. Int. Ass. theoret. et applied Limnology*, XI, 455—458, 1951, 1 portr.

BLUM, J. L. - Notes on Vaucheriaceae, with particular reference to Western New York, *Bull. Torrey Bot. Club*, 78, no 6, 1951, 441—448, 13 fig. Described as new: *Vaucheria brevicaulis* sp. n.

WERFF, A. VAN DER - La croissance des Algues dans les installations néerlandaises d'épuration et de distribution d'eau potable, *Bull. Soc. Bot. Nord de la France*, IV, no 2, 1951, 52—55.

THOMASSON, K. - Beiträge zur Kenntnis des Planktons einiger Seen im nordschwedischen Hochgebirge, *Schweiz. Ztschr. Hydrol.*, XIII/2, 1951, 336—351, 4 fig., 2 tables.

Contribution to the knowledge of the plankton of some lakes of the North-Swedish highlands. With a list of the organisms found.

SYMOENS, J. J. (with P. DUVIGNEAUD, J. DEWIT and C. KIWAK). - Note sur les tufs calcaires de la vallée du Hoyoux, *Lejeunia*, 14, 1950, 13—16, 3 pl., 1 table.

Note on the calcareous tuffs of the Hoyoux-valley.

DUVIGNEAUD, P. & SYMOENS, J. J. - Observations sur la strate algale des formations herbeuses du sud du Congo Belge, *Lejeunia*, 13, 1949, 67—98, 10 pl., 3 tables.

Investigations on the algal stratum of the herbaceous formations in southern Belgian Congo. With description of a new genus: *Cirrosiphon* gen. n. and three new species: *Schizothrix pseudofriesii*, *Sch. luteola* and *Cirrosiphon geniculatus* sp. n.

SYMOENS, J. J. - Quelques acquisitions récentes en Limnologie, I. Quelques données sur la biologie des Lacs, Bruxelles, 1950—51, 97 pp., 21 fig., 7 pl., 3 tables.

Some recent accessions in Limnology, I. Some data on biology of lakes. Four points are discussed: 1. The thermic stratification and its consequences. 2. Plankton. 3. Lake sedimentations, its flora and fauna. 4. Various types of lakes and their evolution.

BUTTIAUX, R. - L'Analyse bactériologique des eaux de consommation,

(Collection de l'Institut Pasteur), Paris, 1951, 209 pp., 38 fig., 21 tables, 1 scheme.

BALDI, E. - Stabilité dans le temps de la biocénose zooplanktique du Lac Maggiore, *Tr. Ass. Int. Limnol. théor. et appl.*, XI, 1951, 35—40, 1 fig., 1 table.

Temporal stability of the zooplankton population in Lago Maggiore. The author thinks that the stability of biocoenotic pattern in a lake, within a few scores of years, is a function of the stability of its ecological conditions and of the hydrographic stability of the lake.

TONOLLI, V. - A new device for continuous plankton sampling: The plankton bar, *Proc. Int. Ass. ther. and appl. Limnology*, XI, 1951, 422—429, 6 fig., 1 table.

RAVERA, O. - Emploi de la Paraldehyde acétique pour la narcose de petits organismes d'eau douce, *Tr. Ass. Int. Limnol. théor. et appl.*, XI, 1951, 315—316, 1 table.

Acetic paraldehyde as a narcotic for small freshwater Invertebrates.

RAVERA, O. - Schéma écologique d'un méandre de rivière alpine, *Tr. Ass. Int. Limnol. théor. et appl.*, XI, 1951, 308—314, 4 fig., 1 table.

Observations on the ecology of a cut-off arm of an alpine river. Preliminary conclusions: 1° The most important factor in the ecology of this environment is the speed of the stream; 2° an interaction between the biological and environmental conditions can be observed; 3° this interaction is determined by the flow of the stream entering the cut-off arm; 4° between the periodically increased flow of the river and the expansion pressure of the benthic biocenoses there is an unstable equilibrium.

DONASZY, E. - Seasonal changes in the dissolved oxygen-concentration of the Szelid-lake Hungary, *Arch. Hydrob.*, XLV, 1951, 314—326, 1 fig., 7 tables, 1 folder.

The following points are discussed at some length: 1° The dissolved oxygen-concentration of the upper layer; 2° The layers of the dissolved oxygen.

BRETSCHNEIDER, L. H. - Elektronenmikroskopische Untersuchung einiger Ziliaten, *Mikroskopie*, 5, 11/12, 1950, 257—269, 10 fig.

Electronic-microscopical research on some Ciliata. *Paramecium caudatum*, *Isotricha ruminantia* and *Opalina dimidiata* were studied; the structure of the cilia, of the ciliate plasm and some other details are described.

STALBERG, G. - Further research into the phytoplankton of the Göta Alv, *Botaniska Notiser*, 1951, H. 3, 241—254, 3 tables, 1 folder.

LENZ, F. - Neue Beobachtungen zur Biologie der Jugendstadien der Tendipedidengattung *Parachironomus* Lenz, *Zool. Anz.*, 147/5/6, 1951, 95—111, 10 fig.

New observations on biology of the early stadia of the Tendipedid genus *Parachironomus* Lenz.

LENZ, F. - Zur Funktion der Tubuli und Analschläuche der Chironomus-larven, *Verh. Dtsch. Zool. Kiel*, 1948, 484—489, 4 fig.  
On the function of the tubuli and of the anal papillae of the Chironomus-larvae.

LENZ, F. - Mittelbare Geschlechter-Findung bei Dipteren, *Verh. Dtsch. Zoologen in Marburg*, 1950, 285—289.  
Indirect sexes-locating in Diptera. Observations on *Phlebotomus papatasi* Scop. (Psychodidae). Author's opinion is that identical instinct and identical physiological structures are the reasons that make both sexes to meet for copulation.

LENZ, F. - Probleme der Chironomiden-Forschung, *Verh. Int. Ver. f. theor. u. angew. Limnologie*, XI, 1951, 230—245.  
The problems of the research on Chironomidae.

JAARCIJFERS OVER DE VISSERIJ GEDURENDE HET JAAR 1950. - *Verslagen en Mededelingen van de Directie van de Visserijen*, no 41, 1950, 108 p., 25 tables.  
Statistics over Fisheries for 1950.

DRESSCHER, TH. G. N. - De invloed van het Amsterdamse Rioolwater op het IJsselmeer vroeger, nu en in de Toekomst, *Ned. Centr. Organisatie voor toegepast natuurwetensch. onderzoek*, 1951, 56 p., 20 tables, 9 graphics.  
The Influence of the Amsterdam Sewage on the IJsselmeer in the past, at present and in the future. With an English summary.

BERICHTE DER DEUTSCHEN WISSENSCHAFTLICHEN KOMMISSION FÜR MEERES-FORSCHUNG, N. F., XII/3, 1951.  
Contents: E. HENTSCHEL, Untersuchungen über das Plankton des Bornholmbeckens. — J. LUNDBECK, Biologisch-statistische Untersuchungen über die deutsche Hochseefischerei.

VISSERIJ-NIEUWS, IV, no 7, Nov. 1951. - Contents: Fishery-results. — Our most important fish-species. — Damage to nets.

BALDI, E. - Phénomènes de microévolution dans les populations planktiques d'eau douce, *Vierteljahrschr. Nat. f. Ges. Zürich*, XCV, 1950, 89—114, 7 fig.  
Microevolutional phenomena in plankton-population in freshwater. The following points are discussed: I. Genetical point of view in Limnology. — II. Plankton-population. — III. Isolation and differentiation in plankton-populations. — IV. Evolution of plankton-populations.

BALDI, E. - Adriano Garbini e il Problema dell'Origine delle Limnofaune, *Mem. Mus. Civ. Storia Nat. Verona*, II, 1950, 333—345.  
Adriano Garbini and the problem of the origins of the limnofaunas

FOX, MUNRO H., - Prof. E. Baldi, - *Nature*, 168, 1951, 496. Obituary.

BALDI, E. - Stabilità delle biocenosi limnetiche e sue probabili condizioni, *Boll. Zool.*, XVIII, 1951, 65—78.  
Stability of the limnetic biocenoses and its probable conditions of it.



- WANDEL, G. - Über den Nutzen und Schaden des Uferbewuchses an fließenden Gewässern, *Arb. Landesamtes f. Gewässerkunde im Min. f. Wirtschaft u. Verkehr des Landes Nordrhein-Westfalen*, 41 p., 63 fig., 31 phot.  
On utility and damage caused by bank-vegetation of running waters.
- BARTOS, E. - Československe druhy rodu *Testudinella* a *Pompholyx* (Rot.), *Sborník Klubu Přírodovědeckého v Brně*, XXVIII, 1951, 10—20, 51 fig.  
Czechoslovakian species of the genera *Testudinella* and *Pompholyx* (Rot.). With an identification-key. Two new species are described, viz. *Testudinella pseudoelliptica* n. sp. and *T. sculpturata* n. sp.
- BARTOS, E. - Korennozci, Virnici a Zelvusky mechu Sumavských predhori, *Casopis Nar. Musea*, 49—64, 2 pl.  
On the Rhizopoda, Rotatoria and Tardigrada from the mosses of the undermountains of Sumava (Bohemia). A new Rhizopod and five new Rotatoria are described, viz. *Chlamydothrys jirovci* n. sp.; *Habrotricha curvicollis* n. sp., *H. incola* n. sp., *H. lamellata* n. sp., *Mniobia discophora* n. sp. and *Philodina amethystina* n. sp.
- BARTOS, E. - Klic k Urcovani Virniku rodu *Polyarthra* Ehrbg., *Kasopis Nar. Musea*, 82—91, 12 fig.  
Key to determination of Rotifera from the genus *Polyarthra* Ehrbg.
- JAHRESBERICHT ÜBER DIE DEUTSCHE FISCHEREI 1950, Berlin 1951.  
Annual Report on German Fishery 1950.
- LAUTERBORN, R. - Zum Geleit, *Arch. f. Hydrobiol.*, 1951, Suppl. Bd. XX, (*Schriften d. hydrobiol. Station f. d. Schwarzwald, Falkau*, I, 1—17).  
An accompanying Word (as introduction to the first series of papers from the Hydrobiological Station at Falkau).

Dr W. JUNK, PUBLISHERS, THE HAGUE, NETHERLANDS

INSECTS AS HUMAN FOOD

by

F. S. BODENHEIMER

Jerusalem

1951. 352 p. w. 75 illustrations . . . . . f 10.—

“The utilization of insects as food was the proper instinctive response to the physiological need for animal proteins, fat or other substances which large quantities of insect food could provide.”

*This is the very book for a gift to your friend.*

PHYSIOLOGIA  
COMPARATA ET  
OECOLOGIA

AN INTERNATIONAL JOURNAL OF  
COMPARATIVE PHYSIOLOGY AND ECOLOGY

EDITED BY:

- |                             |                               |
|-----------------------------|-------------------------------|
| J. TEN CATE, Amsterdam      | CHR. ROMIJN, Utrecht          |
| R. CHAUVIN, Paris           | P. SAWAYA, São Paulo          |
| S. DIJKGRAAF, Utrecht       | P. F. SCHOLANDER, Swartmore   |
| M. FLORKIN, Liège           | E. J. SLIJPER, Amsterdam      |
| M. FONTAINE, Paris          | H. J. VONK, Bogor             |
| H. HEDIGER, Basel           | J. H. WELSH, Cambridge, Mass. |
| B. A. HOUSSAY, Buenos Aires | C. A. G. WIERSMA, Pasadena    |
| C. W. MENG, Peiping         | G. M. YONGE, Glasgow          |
| C. F. A. PANTIN, Cambridge  |                               |

This international quarterly publishes articles dealing with comparative physiology and ecology of Vertebrates and Invertebrates. It also contains contributions on animal biology, for example on nutrition, feeding mechanism, locomotion, reproduction, hibernation, marine biology, etc. The journal contains a limited number of comprehensive book reviews. Text in English, French, and German mixed.

Current issue: Vol. II . . . . . f 36.—



## CONTENTS

COMITA, G. W. Cyclops Magnus from Amchitka, Alaska ....	337
BLANC, M. Etudes consacrées aux animaux dulcicoles et recherches d'Hydrobiologie animale en France de 1945 à 1950 .....	341
MACAN, T. T. Taxonomy of the nymphs of the British species of Leptophlebiidae (Ephem.) .....	363
SCOTT, A. M. and PRESCOTT, G. W. The algal flora of South- eastern United States VI. Additions to our knowledge of the desmid genus Euastrum 2 .....	377
PROSKAUER, J. On the nuclear cytology of Hydrodictyon ....	399
Bibliography .....	409

Prix de souscription d'un volume (env. 400 p. en 4 fasc.) . . fl. holl. 40.—

Subscription price for the volume (about 400 pp. in 4 parts) Dutch fl. 40.—

Abonnement pro Band (ca. 400 Seiten in 4 Heften) . . . Holl. fl. 40.—